

Exact evaluation of density matrix elements for the Heisenberg chain

Jun Sato ¹ ^{*}, Masahiro Shiroishi ¹ [†], Minoru Takahashi ¹ [‡]

¹ *Institute for Solid State Physics, University of Tokyo,
Kashiwanoha 5-1-5, Kashiwa, Chiba 277-8581, Japan*

February 2, 2008

Abstract

We have obtained all the density matrix elements on six lattice sites for the spin-1/2 Heisenberg chain via the algebraic method based on the quantum Knizhnik-Zamolodchikov equations. Several interesting correlation functions, such as chiral correlation functions, dimer-dimer correlation functions, etc... have been analytically evaluated. Furthermore we have calculated all the eigenvalues of the density matrix and analyze the eigenvalue-distribution. As a result the exact von Neumann entropy for the reduced density matrix on six lattice sites has been obtained.

^{*}junji@issp.u-tokyo.ac.jp

[†]siroisi@issp.u-tokyo.ac.jp

[‡]mtaka@issp.u-tokyo.ac.jp

1 Introduction

The spin-1/2 antiferromagnetic Heisenberg XXZ chain is one of the most fundamental models for one-dimensional quantum magnetism, which is given by the Hamiltonian

$$\mathcal{H} = \sum_{j=1}^N (S_j^x S_{j+1}^x + S_j^y S_{j+1}^y + \Delta S_j^z S_{j+1}^z), \quad (1.1)$$

where $S_j^\alpha = \sigma_j^\alpha/2$ with σ_j^α being the Pauli matrices acting on the j -th site, Δ is the anisotropy parameter, and N is the number of lattice sites of this system. Here we impose the periodic boundary condition $\vec{S}_{j+N} = \vec{S}_j$. For $\Delta > 1$, it is called the massive XXZ model where the system is gapful. Meanwhile for $-1 < \Delta \leq 1$ case, the system is gapless and called the massless XXZ model. Especially we call it XXX model for the isotropic case $\Delta = 1$.

The exact eigenvalues and eigenvectors of this model can be obtained by the Bethe Ansatz method [1, 2]. We shall give a brief survey of this method below. First we assume the wave function $|\Psi\rangle$ with M down spins in the form

$$|\Psi\rangle = \sum_{1 \leq n_1 < \dots < n_M \leq N} \sum_{\sigma \in S_n} A(\sigma) \exp \left[i \sum_{j=1}^M k_{\sigma(j)} n_j \right] |n_1, \dots, n_M\rangle, \\ A(\sigma) = \epsilon(\sigma) \prod_{l < j} (e^{i(k_{\sigma(l)} + k_{\sigma(j)})} + 1 - 2\Delta e^{ik_{\sigma(l)}}), \quad (1.2)$$

where σ denotes an element of the symmetric group S_n , $\epsilon(\sigma)$ is the sign of the permutation, and $|n_1, \dots, n_M\rangle$ signifies the state where the spins at the positions $n_1, \dots, n_M$ are downward and all the other spins directing upward. If the quasi-momenta $\{k_1, k_2, \dots, k_M\}$ satisfy the Bethe ansatz equations

$$e^{ik_j N} = \prod_{l \neq j}^M \left(-\frac{e^{i(k_j + k_l)} + 1 - 2\Delta e^{ik_j}}{e^{i(k_j + k_l)} + 1 - 2\Delta e^{ik_l}} \right), \quad (j = 1, 2, \dots, M), \quad (1.3)$$

the wave function (1.2) becomes the eigenvector of the Hamiltonian (1.1):

$$\mathcal{H}|\Psi\rangle = E|\Psi\rangle, \quad E = \frac{N\Delta}{4} + \sum_{j=1}^M (\cos k_j - \Delta). \quad (1.4)$$

In the thermodynamic limit $N \rightarrow \infty$, the ground state energy per site e_0 of the massless XXZ model, for example, is computable by analyzing the Bethe ansatz equations

$$e_0 = \frac{\Delta}{4} - \frac{\sin(\pi\nu)}{2\pi} \int_{-\infty}^{\infty} dt \frac{\sinh[(1-\nu)t]}{\sinh t \cosh(\nu t)}, \quad (1.5)$$

where we parametrize the anisotropy parameter as $\Delta = \cos(\pi\nu)$. Especially in the XXX case $\Delta = 1$, the ground state energy per site (1.5) can be simplified into

$$e_0 = \frac{1}{4} - \ln 2. \quad (1.6)$$

In the same way other physical quantities in the thermodynamic limit such as specific heat, magnetic susceptibility, elementary excitations, etc..., can be exactly evaluated even at finite temperature by the Bethe ansatz method [2].

The exact calculation of the correlation functions, however, is still a difficult problem even in the simplest case for static correlation functions at zero temperature. The exceptional case is $\Delta = 0$, where the system reduces to a lattice free-fermion model by the Jordan-Wigner transformation. In this case, we can calculate arbitrary correlation functions by means of Wick's theorem [3, 4]. However, there have been rapid developments recently in the exact evaluations of correlation functions for $\Delta \neq 0$ case also. Below we shall give the historical review of them, concentrating mainly on the static correlation functions of the XXX model $\Delta = 1$ at zero temperature for the infinite system $N \rightarrow \infty$. Until quite recently only the correlation functions within three lattice sites had been known:

$$\langle S_1^z S_2^z \rangle = \frac{1}{12} - \frac{1}{3} \zeta_a(1) = -0.1477157268533151 \dots, \quad (1.7)$$

$$\langle S_1^z S_3^z \rangle = \frac{1}{12} - \frac{4}{3} \zeta_a(1) + \zeta_a(3) = 0.06067976995643530 \dots, \quad (1.8)$$

where $\zeta_a(s)$ is the alternating zeta function defined by $\zeta_a(s) \equiv \sum_{n=1}^{\infty} (-1)^{n-1} / n^s$, which is related to Riemann zeta function $\zeta(s) \equiv \sum_{n=1}^{\infty} 1/n^s$ as $\zeta_a(s) = (1 - 2^{1-s})\zeta(s)$. Note that the alternating zeta function is regular at $s = 1$ and is given by $\zeta_a(1) = \ln 2$. The nearest-neighbor correlation function (1.7) is derived directly from the ground state energy (1.6) obtained by Hulthén in 1938 [5]. The first nontrivial correlation function (1.8) was derived by Takahashi in 1977 via the strong coupling expansion of the ground state energy for the half-filled Hubbard model [6]. Note also that another derivation of the second-neighbor correlation function (1.8) was given by Dittrich and Inozemtsev in 1997 [7]. These method, however, can not be generalized to calculate further correlation functions on more than four lattice sites, unfortunately.

Meanwhile using the representation theory of the quantum affine algebra $U_q(\hat{sl}_2)$, Kyoto Group (Jimbo, Miki, Miwa, Nakayashiki) derived a multiple integral representation for arbitrary correlation functions of the massive XXZ antiferromagnetic chain in 1992 [8, 9], which is before long extended to the XXX case [10, 11] and the massless XXZ case [12]. More recently the same integral representations were reproduced by Kitanine, Maillet, Terras [13] in the framework of Quantum Inverse Scattering Method. They have also succeeded in generalizing the integral representations to the XXZ model with an external magnetic field [13]. More recently the integral formulas are extended to the XXZ model even at finite temperature ! [14, 15]. In this way it has been established now the correlation functions on n lattice sites are represented by n -dimensional integrals in general. However, these multiple integral representations for correlation functions, though completely exact, have not been used widely especially among physicists. It is mainly because we can not evaluate accurate numerical values of correlation functions directly from the integral representation. Also it had been a puzzle that the exact expression of the correlation function on three lattice sites (1.8) had not been reproduced from the integral representation for a long time.

The situation changed when Boos and Korepin devised an innovative method to evaluate these multiple integrals for XXX chain in 2001 [16, 17]. They showed that the integrand in the multiple integral formula can be reduced to a *canonical* form, which allows us to implement the integration. This method was at first applied to a special correlation function called Emptiness Formation Probability (EFP) [11] which signifies the probability to find a

ferromagnetic string of length n :

$$P(n) \equiv \left\langle \prod_{j=1}^n \left(\frac{1}{2} + S_j^z \right) \right\rangle. \quad (1.9)$$

By performing the multiple integrals, the explicit forms of the EFP for up to $n = 5$ was obtained [16, 17, 18].

The Boos-Korepin method was applied to calculate other correlation functions on four lattice sites in 2003 [19]. Especially the third-neighbor correlation function was obtained there as

$$\begin{aligned} \langle S_1^z S_4^z \rangle &= \frac{1}{12} - 3\zeta_a(1) + \frac{74}{9}\zeta_a(3) - \frac{56}{9}\zeta_a(1)\zeta_a(3) - \frac{8}{3}\zeta_a(3)^2 - \frac{50}{9}\zeta_a(5) + \frac{80}{9}\zeta_a(1)\zeta_a(5) \\ &= -0.05024862725723524 \dots \end{aligned} \quad (1.10)$$

Other correlation functions on four lattice sites are given in Appendix A.

Subsequently, the Boos-Korepin method was generalized to XXZ chain both in massless and massive regime and all the correlation functions within four lattice sites were obtained for general anisotropy [20, 21, 22].

In principle, multiple integrals for any correlation functions can be performed by means of Boos-Korepin method. However, $P(5)$ is the only correlation function which was calculated by this method on five lattice sites, since it is getting critically harder to reduce the integrand to canonical form as the integral dimension increases.

In the course of attempting to obtain further correlation functions, the alternative algebraic method to calculate the EFP was developed by Boos, Korepin and Smirnov in 2003 [23]. They considered the inhomogeneous XXX model, in which each site carries an inhomogeneous parameter λ_j . The homogeneous XXX model corresponds to the case with all the inhomogeneous parameters λ_j set to be 0. Inhomogeneous correlation functions on n lattice sites are considered to be functions of variables $\lambda_1, \dots, \lambda_n$. They derived the functional relations which the inhomogeneous EFP should satisfy by investigating the quantum Knizhnik-Zamolodchikov (qKZ) equations [24, 25, 26, 27], the solutions to which are connected with the inhomogeneous correlation functions [9, 10, 12]. Moreover they suggested an ansatz for the form of the inhomogeneous EFP, which consists of only one transcendental function $\omega(\lambda)$ with rational functions of inhomogeneous parameters $\lambda_1, \dots, \lambda_n$ as coefficients. (for the proof of the ansatz and further generalizations, see [28, 29, 30, 31, 32, 33]). It was shown that the functional relations together with the ansatz for the final form completely fix the explicit form of the inhomogeneous EFP. In this way the explicit forms of the inhomogeneous EFP for up to $n = 6$ have been obtained, which gives a new result for $P(6)$ in the homogeneous limit $\lambda_j \rightarrow 0$.

This Boos-Korepin-Smirnov method based on the qKZ equation was generalized to arbitrary correlation functions in [34]. Actually, in that paper, all the correlation functions on five lattice sites are obtained based on the functional relations for the general correlation functions (see Appendix A). Especially the fourth-neighbor correlation function is given by

$$\begin{aligned} \langle S_1^z S_5^z \rangle &= \frac{1}{12} - \frac{16}{3}\zeta_a(1) + \frac{290}{9}\zeta_a(3) - 72\zeta_a(1)\zeta_a(3) - \frac{1172}{9}\zeta_a(3)^2 - \frac{700}{9}\zeta_a(5) \\ &\quad + \frac{4640}{9}\zeta_a(1)\zeta_a(5) - \frac{220}{9}\zeta_a(3)\zeta_a(5) - \frac{400}{3}\zeta_a(5)^2 + \frac{455}{9}\zeta_a(7) - \frac{3920}{9}\zeta_a(1)\zeta_a(7) \\ &\quad + 280\zeta_a(3)\zeta_a(7) = 0.03465277698272816 \dots \end{aligned} \quad (1.11)$$

The main purpose of this paper is to report further results using the algebraic method in [34]. More explicitly we have succeeded in calculating all the correlation functions on six lattice sites using this algebraic method. We remark that, if we consider only the two-point correlation functions $\langle S_1^z S_n^z \rangle$, it was already calculated up to the seventh-neighbor correlation function $\langle S_1^z S_8^z \rangle$ [35, 36]. The method actually allows us to evaluate some other correlation functions such as string correlation functions [37], but not all the correlation functions. We also remark that there is a related but different approach for the evaluation of the general correlation functions, which is developed recently by Boos, Jimbo, Miwa, Smirnov and Takeyama [28, 29, 30, 31, 32, 33]. They have established a compact exponential formula for general correlation functions without heavy multiple integrals, which would be also useful to evaluate correlation functions analytically. We, however, do not discuss the formula in this paper.

The plan of this paper is as follows. In section 2, the algebraic method to calculate general correlation functions is reviewed. In section 3, we present the analytical results for some physically interesting correlation functions, such as chiral correlation functions, dimer-dimer correlation functions, etc. . . . In section 4, we show the eigenvalue-distribution of the reduced density matrix and calculate the von Neumann entropy. Summary and discussion are given in section 5.

2 Algebraic method to evaluate general correlation functions based on the qKZ equations

Below we describe the functional approach to evaluate general correlation functions established in [34]. Any correlation function can be expressed as a sum of density matrix elements $P_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}$, which are defined by the ground state expectation value of the product of elementary matrices:

$$P_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n} \equiv \langle E_1^{\epsilon'_1 \epsilon_1} \dots E_n^{\epsilon'_n \epsilon_n} \rangle, \quad (2.1)$$

where $E_j^{\epsilon'_j \epsilon_j}$ are 2×2 elementary matrices acting on the j -th site as

$$\begin{aligned} E_j^{++} &= \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}_{[j]} = \frac{1}{2} + S_j^z, & E_j^{--} &= \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}_{[j]} = \frac{1}{2} - S_j^z, \\ E_j^{+-} &= \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}_{[j]} = S_j^+ = S_j^x + iS_j^y, & E_j^{-+} &= \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}_{[j]} = S_j^- = S_j^x - iS_j^y. \end{aligned}$$

For example, the density matrix elements on four lattice sites P_{++--}^{-++-} can be written in terms of spin operators as

$$P_{++--}^{-++-} = \left\langle S_1^- \left(\frac{1}{2} + S_2^z \right) S_3^+ \left(\frac{1}{2} - S_4^z \right) \right\rangle. \quad (2.2)$$

First we rewrite the functional relations for inhomogeneous density matrix elements, which follow from the qKZ equations:

- Translational invariance

$$P_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(\lambda_1 + \lambda, \dots, \lambda_n + \lambda) = P_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(\lambda_1, \dots, \lambda_n), \quad (2.3)$$

- Transposition, Negating and Reverse order relations

$$\begin{aligned} P_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(\lambda_1, \dots, \lambda_n) &= P_{\epsilon'_1, \dots, \epsilon'_n}^{\epsilon_1, \dots, \epsilon_n}(-\lambda_1, \dots, -\lambda_n) \\ &= P_{-\epsilon_1, \dots, -\epsilon_n}^{-\epsilon'_1, \dots, -\epsilon'_n}(\lambda_1, \dots, \lambda_n) = P_{\epsilon_n, \dots, \epsilon_1}^{\epsilon'_n, \dots, \epsilon'_1}(-\lambda_n, \dots, -\lambda_1) \end{aligned} \quad (2.4)$$

- Intertwining relation

$$\begin{aligned} &\sum_{\tilde{\epsilon}'_j, \tilde{\epsilon}'_{j+1} = \pm} R_{\tilde{\epsilon}'_j \tilde{\epsilon}'_{j+1}}^{\epsilon'_j \epsilon'_{j+1}}(\lambda_j - \lambda_{j+1}) P_{\dots, \epsilon_{j+1}, \epsilon_j, \dots}^{\dots, \tilde{\epsilon}'_{j+1}, \tilde{\epsilon}'_j, \dots}(\dots, \lambda_{j+1}, \lambda_j, \dots) \\ &= \sum_{\tilde{\epsilon}_j, \tilde{\epsilon}_{j+1} = \pm} P_{\dots, \tilde{\epsilon}_j, \tilde{\epsilon}_{j+1}, \dots}^{\dots, \epsilon'_j, \epsilon'_{j+1}, \dots}(\dots, \lambda_j, \lambda_{j+1}, \dots) R_{\epsilon_j \epsilon_{j+1}}^{\tilde{\epsilon}_j \tilde{\epsilon}_{j+1}}(\lambda_j - \lambda_{j+1}), \end{aligned} \quad (2.5)$$

where R denotes the R -matrix of the XXX model:

$$R(\lambda) = \begin{pmatrix} R_{++}^{++}(\lambda) & R_{+-}^{++}(\lambda) & R_{-+}^{++}(\lambda) & R_{--}^{++}(\lambda) \\ R_{++}^{+-}(\lambda) & R_{+-}^{+-}(\lambda) & R_{-+}^{+-}(\lambda) & R_{--}^{+-}(\lambda) \\ R_{++}^{-+}(\lambda) & R_{+-}^{-+}(\lambda) & R_{-+}^{-+}(\lambda) & R_{--}^{-+}(\lambda) \\ R_{++}^{--}(\lambda) & R_{+-}^{--}(\lambda) & R_{-+}^{--}(\lambda) & R_{--}^{--}(\lambda) \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{\lambda}{\lambda+1} & \frac{1}{\lambda+1} & 0 \\ 0 & \frac{1}{\lambda+1} & \frac{\lambda}{\lambda+1} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (2.6)$$

- First recurrent relation

$$\begin{aligned} P_{\epsilon_1 \epsilon_2, \dots, \epsilon_n}^{\epsilon'_1 \epsilon'_2, \dots, \epsilon'_n}(\lambda + 1, \lambda, \lambda_3, \dots, \lambda_n) &= -\delta_{\epsilon_1, -\epsilon_2} \epsilon'_1 \epsilon'_2 P_{-\epsilon'_1 \epsilon'_3, \dots, \epsilon'_n}^{\epsilon'_2 \epsilon'_3, \dots, \epsilon'_n}(\lambda, \lambda_3, \dots, \lambda_n) \\ P_{\epsilon_1 \epsilon_2, \dots, \epsilon_n}^{\epsilon'_1 \epsilon'_2, \dots, \epsilon'_n}(\lambda - 1, \lambda, \lambda_3, \dots, \lambda_n) &= -\delta_{\epsilon'_1, -\epsilon'_2} \epsilon'_1 \epsilon'_2 P_{\epsilon_2 \epsilon_3, \dots, \epsilon_n}^{-\epsilon'_1 \epsilon'_3, \dots, \epsilon'_n}(\lambda, \lambda_3, \dots, \lambda_n) \end{aligned} \quad (2.7)$$

- Second recurrent relation

$$\lim_{\lambda_j \rightarrow i\infty} P_{\epsilon_1, \dots, \epsilon_j, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_j, \dots, \epsilon'_n}(\lambda_1, \dots, \lambda_j, \dots, \lambda_n) = \delta_{\epsilon_j, \epsilon'_j} \frac{1}{2} P_{\epsilon_1, \dots, \epsilon_{j-1}, \epsilon_{j+1}, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_{j-1}, \epsilon'_{j+1}, \dots, \epsilon'_n}(\lambda_1, \dots, \lambda_{j-1}, \lambda_{j+1}, \dots, \lambda_n) \quad (2.8)$$

- Identity relation

$$\begin{aligned} \sum_{\epsilon_1, \dots, \epsilon_n} P_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(\lambda_1, \dots, \lambda_n) &= \sum_{\epsilon'_1, \dots, \epsilon'_n} P_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(\lambda_1, \dots, \lambda_n) \\ &= P_{+, \dots, +}^{+, \dots, +}(\lambda_1, \dots, \lambda_n) = P_{-, \dots, -}^{-, \dots, -}(\lambda_1, \dots, \lambda_n) \end{aligned} \quad (2.9)$$

- Reduction relation

$$\begin{aligned} P_{+, \epsilon_2, \dots, \epsilon_n}^{+, \epsilon'_2, \dots, \epsilon'_n}(\lambda_1, \lambda_2, \dots, \lambda_n) + P_{-, \epsilon_2, \dots, \epsilon_n}^{-, \epsilon'_2, \dots, \epsilon'_n}(\lambda_1, \lambda_2, \dots, \lambda_n) &= P_{\epsilon_2, \dots, \epsilon_n}^{\epsilon'_2, \dots, \epsilon'_n}(\lambda_2, \dots, \lambda_n) \\ P_{\epsilon_1, \dots, \epsilon_{n-1}, +}^{\epsilon'_1, \dots, \epsilon'_{n-1}, +}(\lambda_1, \dots, \lambda_{n-1}, \lambda_n) + P_{\epsilon_1, \dots, \epsilon_{n-1}, -}^{\epsilon'_1, \dots, \epsilon'_{n-1}, -}(\lambda_1, \dots, \lambda_{n-1}, \lambda_n) &= P_{\epsilon_1, \dots, \epsilon_{n-1}}^{\epsilon'_1, \dots, \epsilon'_{n-1}}(\lambda_1, \dots, \lambda_{n-1}) \end{aligned} \quad (2.10)$$

Additionally it is established that density matrix elements can be written in the form of [23, 28, 34]

$$P_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(\lambda_1, \dots, \lambda_n) = \left(\prod_{j=1}^n \frac{\delta_{\epsilon_j, \epsilon'_j}}{2} \right) + \sum_{l=1}^{\lfloor \frac{n}{2} \rfloor} \sum_{\sigma \in T_{n,l}} A_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(l, \sigma | \lambda_{\sigma(1)}, \dots, \lambda_{\sigma(n)}) \prod_{j=1}^l \omega(\lambda_{\sigma(2j-1)} - \lambda_{\sigma(2j)}), \quad (2.11)$$

where $T_{n,l}$ is a subset of the symmetry group of degree n defined as

$$T_{n,l} = \{ \sigma \in S_n | \sigma(1) < \sigma(3) < \dots < \sigma(2l-1), \\ \sigma(2j-1) < \sigma(2j) \quad \text{for } j = 1, 2, \dots, l, \\ \sigma(2l+1) < \sigma(2l+2) < \dots < \sigma(n) \}. \quad (2.12)$$

$A_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(l, \sigma | \lambda_1, \dots, \lambda_n)$ are rational functions of inhomogeneous parameters $\lambda_1, \dots, \lambda_n$ with known denominator

$$A_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(l, \sigma | \lambda_1, \dots, \lambda_n) = \frac{Q_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(l, \sigma | \lambda_1, \dots, \lambda_n)}{D_{n,l}(\lambda_1, \dots, \lambda_n)}, \\ D_{n,l}(\lambda_1, \dots, \lambda_n) = \frac{\prod_{k=1}^l (\lambda_{2k-1} - \lambda_{2k}) \prod_{2l+1 \leq k < j \leq n} (\lambda_k - \lambda_j)}{\prod_{1 \leq k < j \leq n} (\lambda_k - \lambda_j)}. \quad (2.13)$$

$Q_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(l, \sigma | \lambda_1, \dots, \lambda_n)$ are polynomials of known degree in inhomogeneous parameters $\lambda_1, \dots, \lambda_n$ with rational coefficients which have at most the same degree in each variable and also at most the same total degree as in the denominator.

A transcendental function $\omega(\lambda)$ in (2.11) is defined by

$$\omega(\lambda) = \frac{1}{2} + \sum_{k=1}^{\infty} (-1)^k \frac{2k(\lambda^2 - 1)}{\lambda^2 - k^2} = 2 \sum_{k=0}^{\infty} \lambda^{2k} \{ \zeta_a(2k-1) - \zeta_a(2k+1) \}. \quad (2.14)$$

Here note that the identity $\zeta_a(-1) = -3\zeta(-1) = 1/4$. The following properties of the function $\omega(\lambda)$ are needed for calculations of the recurrent relations (2.7)-(2.8):

$$\omega(i\infty) = 0, \quad \omega(\lambda \pm 1) = \frac{3}{2} \frac{1}{\lambda^2 - 1} - \frac{\lambda(\lambda \pm 2)}{\lambda^2 - 1} \omega(\lambda). \quad (2.15)$$

Below we list the explicit forms of all the non-zero elements of inhomogeneous density matrices for $n = 1, 2$:

$$P_+^+(\lambda_1) = P_-^-(\lambda_1) = \frac{1}{2}, \\ P_{++}^{++}(\lambda_1, \lambda_2) = P_{--}^{--}(\lambda_1, \lambda_2) = \frac{1}{4} + \frac{1}{6} \omega(\lambda_1 - \lambda_2), \\ P_{+-}^{+-}(\lambda_1, \lambda_2) = P_{-+}^{-+}(\lambda_1, \lambda_2) = \frac{1}{4} - \frac{1}{6} \omega(\lambda_1 - \lambda_2), \\ P_{-+}^{+-}(\lambda_1, \lambda_2) = P_{+-}^{-+}(\lambda_1, \lambda_2) = \frac{1}{3} \omega(\lambda_1 - \lambda_2). \quad (2.16)$$

It can be easily seen that these satisfy the functional relations (2.3)-(2.10). By taking the homogeneous limit $\lambda_1, \lambda_2 \rightarrow 0$, we obtain density matrix elements for the physically interesting homogeneous case:

$$\begin{aligned} P_+^+ &= P_-^- = \frac{1}{2}, & P_{++}^{++} &= P_{--}^{--} = \frac{1}{3} - \frac{1}{3}\zeta_a(1), \\ P_{+-}^{+-} &= P_{-+}^{-+} = \frac{1}{6} + \frac{1}{3}\zeta_a(1), & P_{-+}^{+-} &= P_{+-}^{-+} = \frac{1}{6} - \frac{2}{3}\zeta_a(1). \end{aligned} \quad (2.17)$$

In the paper [34], it has been argued that the functional relations (2.3)-(2.10) together with the ansatz for the form of inhomogeneous correlation functions (2.11) completely determine the unknown rational coefficients in the polynomials $Q_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(l, \sigma | \lambda_1, \dots, \lambda_n)$ and therefore the full form of the inhomogeneous correlation functions. As an example, we give the explicit form of the density matrix element for $n = 3$

$$\begin{aligned} Q_{++-}^{-++}(l=1, \sigma=(123) | \lambda_1, \lambda_2, \lambda_3) &= \frac{1}{6} - \frac{1}{12}\lambda_{12}, \\ Q_{++-}^{-++}(l=1, \sigma=(231) | \lambda_1, \lambda_2, \lambda_3) &= \frac{1}{6} + \frac{1}{12}\lambda_{12}, \\ Q_{++-}^{-++}(l=1, \sigma=(132) | \lambda_1, \lambda_2, \lambda_3) &= \frac{1}{6} + \frac{1}{6}\lambda_{13}\lambda_{23} - \frac{1}{12}(\lambda_{13} + \lambda_{23}), \\ P_{++-}^{-++}(\lambda_1, \lambda_2, \lambda_3) &= \frac{2 - \lambda_{12}}{12\lambda_{13}\lambda_{23}}\omega_{12} + \frac{2 + \lambda_{23}}{12\lambda_{21}\lambda_{31}}\omega_{23} + \frac{2 + 2\lambda_{12}\lambda_{32} - (\lambda_{12} + \lambda_{32})}{12\lambda_{12}\lambda_{32}}\omega_{13}, \end{aligned} \quad (2.18)$$

where we have used the abbreviations $\omega_{jk} = \omega(\lambda_{jk})$, $\lambda_{jk} = \lambda_j - \lambda_k$. By taking the homogeneous limit $\lambda_1, \lambda_2, \lambda_3 \rightarrow 0$, we obtain the homogeneous density matrix element

$$P_{++-}^{-++} = \frac{1}{12} - \frac{4}{3}\zeta_a(1) + \zeta_a(3). \quad (2.19)$$

In this way all the correlation functions on five lattice sites have been obtained in [34]. In this paper we have obtained further results for *six* lattice sites. There are 24 independent correlation functions among them, explicit forms of which are collected in Appendix B.

Concerning two-point correlation functions $\langle S_1^z S_n^z \rangle$, on the other hand, an efficient way to calculate them has been developed [35, 36]. Namely, without evaluating all the density matrix elements, a two-point correlation function can be derived from its generating function

$$P_n^\kappa \equiv \left\langle \prod_{j=1}^n \left\{ \left(\frac{1}{2} + S_j^z \right) + \kappa \left(\frac{1}{2} - S_j^z \right) \right\} \right\rangle \quad (2.20)$$

through the relation

$$\langle S_1^z S_n^z \rangle = \frac{1}{2} \frac{\partial^2}{\partial \kappa^2} \left\{ P_n^\kappa - 2P_{n-1}^\kappa + P_{n-2}^\kappa \right\} \bigg|_{\kappa=1} - \frac{1}{4}. \quad (2.21)$$

Note that the generating function P_n^κ can be considered as a generalization of the Emptiness Formation Probability (1.9), which is reproduced if we set $\kappa = 0$ in (2.20). From the functional relations for general density matrix elements (2.3)-(2.10), it can be shown that the generating function P_n^κ satisfies the following closed functional relations:

- Translational invariance

$$P_n^\kappa(\lambda_1 + \lambda, \dots, \lambda_n + \lambda) = P_n^\kappa(\lambda_1, \dots, \lambda_n) \quad (2.22)$$

- Negating relation

$$P_n^\kappa(-\lambda_1, \dots, -\lambda_n) = P_n^\kappa(\lambda_1, \dots, \lambda_n) \quad (2.23)$$

- Symmetry relation

$$P_n^\kappa(\lambda_1, \dots, \lambda_n) = P_n^\kappa(\lambda_{\sigma(1)}, \dots, \lambda_{\sigma(n)}), \quad (2.24)$$

where σ denotes any element of the symmetry group S_n .

- First recurrent relation

$$P_n^\kappa(\lambda_1, \dots, \lambda_{n-1}, \lambda_{n-1} \pm 1) = \kappa P_{n-2}^\kappa(\lambda_1, \dots, \lambda_{n-2}) \quad (2.25)$$

- Second recurrent relation

$$\lim_{\lambda_n \rightarrow i\infty} P_n^\kappa(\lambda_1, \dots, \lambda_{n-1}, \lambda_n) = \frac{1 + \kappa}{2} P_{n-1}^\kappa(\lambda_1, \dots, \lambda_{n-1}) \quad (2.26)$$

One can explicitly calculate the generating functions P_n^κ recursively with respect to n from these functional relations together with the ansatz for the final form

$$P_n^\kappa(\lambda_1, \dots, \lambda_n) = \left(\frac{1 + \kappa}{2} \right)^n + \sum_{l=1}^{\lfloor \frac{n}{2} \rfloor} \sum_{\sigma \in T_{n,l}} A_{n,l}^\kappa(\lambda_{\sigma(1)}, \dots, \lambda_{\sigma(n)}) \prod_{j=1}^l \omega(\lambda_{\sigma(2j-1)} - \lambda_{\sigma(2j)}),$$

$$A_{n,l}^\kappa(\lambda_1, \dots, \lambda_n) = \frac{Q_{n,l}^\kappa(\lambda_1, \dots, \lambda_n)}{D_{n,l}(\lambda_1, \dots, \lambda_n)}, \quad (2.27)$$

where $T_{n,l}$, $D_{n,l}(\lambda_1, \dots, \lambda_n)$ and $\omega(\lambda)$ are the same as in (2.11) and $Q_{n,l}^\kappa(\lambda_1, \dots, \lambda_n)$ are polynomials containing the parameter κ . The great advantage of this method is that the polynomials $Q_{n,l}^\kappa(\lambda_1, \dots, \lambda_n)$ do not depend on the permutation $\sigma \in T_{n,l}$ due to the symmetry relation (2.24), unlike the case of general density matrix elements $Q_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n}(l, \sigma | \lambda_1, \dots, \lambda_n)$. This fact considerably reduces the amount of bothersome calculation. Actually by using this method, up to the seventh-neighbor correlation function $\langle S_1^z S_8^z \rangle$ have been calculated [36]. Calculating all the density matrix elements is much harder. We, therefore, have succeeded only up to six sites yet.

3 Applications to physically interesting correlation functions

In this section, we discuss several physically interesting correlation functions, which can be evaluated exactly from our results of all the density matrix elements for $n = 6$.

3.1 Chiral correlation function

First let us consider the vector chiral correlation function

$$\left\langle \left(\vec{S}_1 \times \vec{S}_2 \right) \cdot \left(\vec{S}_j \times \vec{S}_{j+1} \right) \right\rangle = 6 \left(\langle S_1^x S_2^z S_j^x S_{j+1}^z \rangle - \langle S_1^x S_2^z S_j^z S_{j+1}^x \rangle \right), \quad (3.1)$$

which measures a chirality of the spin alignment. It has been observed that they have simple expressions [34, 38] compared with the other correlation functions on the same lattice sites

$$\left\langle \left(\vec{S}_1 \times \vec{S}_2 \right) \cdot \left(\vec{S}_3 \times \vec{S}_4 \right) \right\rangle = \frac{1}{2} (\zeta_a(1) - \zeta_a(3)) = -0.1041977484048752 \dots, \quad (3.2)$$

$$\left\langle \left(\vec{S}_1 \times \vec{S}_2 \right) \cdot \left(\vec{S}_4 \times \vec{S}_5 \right) \right\rangle = \frac{1}{2} (\zeta_a(1) - \zeta_a(3)) - \frac{5}{4} (\zeta_a(3) - \zeta_a(5)) = -0.01597638205835821 \dots. \quad (3.3)$$

Here we have newly obtained the vector chiral correlation function for six lattice sites, which has also a simple form though it contains some quadratic terms

$$\begin{aligned} \left\langle \left(\vec{S}_1 \times \vec{S}_2 \right) \cdot \left(\vec{S}_5 \times \vec{S}_6 \right) \right\rangle &= \frac{1}{2} \zeta_a(1) - \frac{11}{3} \zeta_a(3) + 9 \zeta_a(5) - \frac{35}{6} \zeta_a(7) + \frac{4}{3} \zeta_a(1) \zeta_a(3) \\ &\quad + \frac{4}{3} \zeta_a(3)^2 - \frac{32}{3} \zeta_a(1) \zeta_a(5) - \frac{4}{3} \zeta_a(3) \zeta_a(5) + \frac{28}{3} \zeta_a(1) \zeta_a(7) \\ &= -0.01774606473688137 \dots. \end{aligned} \quad (3.4)$$

It is interesting to note that the numerical values of these are all negative and their absolute values are oscillating.

Furthermore we obtain the scalar chiral correlation function defined as below, which reveals an intriguing factorization:

$$\begin{aligned} &\left\langle \left[\left(\vec{S}_1 \times \vec{S}_2 \right) \cdot \vec{S}_3 \right] \left[\left(\vec{S}_4 \times \vec{S}_5 \right) \cdot \vec{S}_6 \right] \right\rangle \\ &= 6 \left(\langle S_1^x S_2^y S_3^z S_4^x S_5^y S_6^z \rangle + \langle S_1^x S_2^y S_3^z S_4^y S_5^z S_6^x \rangle + \langle S_1^x S_2^y S_3^z S_4^z S_5^x S_6^y \rangle \right. \\ &\quad \left. - \langle S_1^x S_2^y S_3^z S_4^z S_5^y S_6^x \rangle - 2 \langle S_1^x S_2^y S_3^z S_4^x S_5^z S_6^y \rangle \right) \\ &= \frac{7}{6} (\zeta_a(1) - 1/4) (\zeta_a(7) - \zeta_a(5)) - \frac{1}{6} (\zeta_a(3) - \zeta_a(1)) (\zeta_a(5) - \zeta_a(3)) \\ &= 0.008133862120680087 \dots. \end{aligned} \quad (3.5)$$

Note that the numerical value in this case is positive. These numerical values may indicate a classical picture of the spin alignment for the antiferromagnetic ground state of quantum Heisenberg chain.

3.2 One-particle Green function

By the Jordan-Wigner transformation

$$S_j^+ = \prod_{k=1}^{j-1} (1 - 2c_k^\dagger c_k) c_j, \quad S_j^- = \prod_{k=1}^{j-1} (1 - 2c_k^\dagger c_k) c_j^\dagger, \quad (3.6)$$

the XXX model is transformed into the isotropic spinless fermion model. Let us consider the one-particle Green function $\langle c_1^\dagger c_n \rangle_f$ for this model. Here the bracket $\langle \dots \rangle_f$ means the

expectation value in the half-filled state of the spinless fermion model. The first-neighbor one-particle Green function $\langle c_1^\dagger c_2 \rangle_f$ is directly obtained from the Hulthen's result (1.7) as

$$\langle c_1^\dagger c_2 \rangle_f = \langle S_1^+ S_2^- \rangle = 2\langle S_1^z S_2^z \rangle = \frac{1}{6} - \frac{2}{3}\zeta_a(1) = -0.2954314537066302 \dots \quad (3.7)$$

The first non-trivial result for this one-particle Green function has been obtained in [19]

$$\begin{aligned} \langle c_1^\dagger c_4 \rangle_f &= 4\langle S_1^+ S_2^z S_3^z S_4^- \rangle = 8\langle S_1^x S_2^z S_3^z S_4^x \rangle \\ &= \frac{1}{30} - 2\zeta_a(1) + \frac{338}{45}\zeta_a(3) - \frac{40}{9}\zeta_a(1)\zeta_a(3) - \frac{32}{15}\zeta_a(3)^2 - \frac{52}{9}\zeta_a(5) + \frac{64}{9}\zeta_a(1)\zeta_a(5) \\ &= 0.08228771668643698 \dots \end{aligned} \quad (3.8)$$

because $\langle c_1^\dagger c_n \rangle_f = 0$ if n is odd. Here we have newly obtained the fifth-neighbor one-particle Green function

$$\begin{aligned} \langle c_1^\dagger c_6 \rangle_f &= 16\langle S_1^+ S_2^z S_3^z S_4^z S_5^z S_6^- \rangle = 32\langle S_1^x S_2^z S_3^z S_4^z S_5^z S_6^x \rangle \\ &= \frac{1}{70} - \frac{10}{3}\zeta_a(1) + 74\zeta_a(3) - \frac{3608}{9}\zeta_a(1)\zeta_a(3) - \frac{90832}{45}\zeta_a(3)^2 - \frac{100288}{135}\zeta_a(3)^3 \\ &\quad - \frac{207464}{315}\zeta_a(5) + \frac{133456}{15}\zeta_a(1)\zeta_a(5) + \frac{3088}{9}\zeta_a(3)\zeta_a(5) + \frac{200576}{45}\zeta_a(1)\zeta_a(3)\zeta_a(5) \\ &\quad - \frac{60704}{45}\zeta_a(3)^2\zeta_a(5) - \frac{1943840}{63}\zeta_a(5)^2 + \frac{242816}{9}\zeta_a(1)\zeta_a(5)^2 + \frac{46112}{9}\zeta_a(3)\zeta_a(5)^2 \\ &\quad + \frac{490880}{189}\zeta_a(5)^3 + \frac{89918}{45}\zeta_a(7) - \frac{308392}{9}\zeta_a(1)\zeta_a(7) + \frac{623128}{9}\zeta_a(3)\zeta_a(7) \\ &\quad - \frac{424928}{9}\zeta_a(1)\zeta_a(3)\zeta_a(7) - \frac{645568}{45}\zeta_a(3)^2\zeta_a(7) + \frac{89536}{9}\zeta_a(5)\zeta_a(7) \\ &\quad - \frac{98176}{9}\zeta_a(3)\zeta_a(5)\zeta_a(7) - \frac{236432}{9}\zeta_a(7)^2 + \frac{343616}{9}\zeta_a(1)\zeta_a(7)^2 - \frac{7052}{5}\zeta_a(9) \\ &\quad + \frac{128928}{5}\zeta_a(1)\zeta_a(9) - \frac{196304}{3}\zeta_a(3)\zeta_a(9) + \frac{645568}{15}\zeta_a(1)\zeta_a(3)\zeta_a(9) \\ &\quad + \frac{98176}{5}\zeta_a(3)^2\zeta_a(9) + \frac{135104}{3}\zeta_a(5)\zeta_a(9) - \frac{196352}{3}\zeta_a(1)\zeta_a(5)\zeta_a(9) \\ &= -0.04497471675792834 \dots \end{aligned} \quad (3.9)$$

It will be an interesting problem to study the asymptotic behavior of one-particle Green function in detail. Conformal field theory predict that

$$\langle c_1^\dagger c_x \rangle_f \sim \cos(k_F x) x^{-5/4}, \quad k_F = \pi/2, \quad (3.10)$$

omitting the logarithmic correction [39]. Unfortunately our new exact results are not sufficient to confirm the asymptotic formula.

3.3 Dimer-dimer correlation function

Next let us consider the dimer-dimer correlation function defined as

$$D_n \equiv \left\langle \left(\vec{S}_1 \cdot \vec{S}_2 \right) \left(\vec{S}_n \cdot \vec{S}_{n+1} \right) \right\rangle = 3\langle S_1^z S_2^z S_n^z S_{n+1}^z \rangle + 6\langle S_1^x S_2^x S_n^z S_{n+1}^z \rangle, \quad (3.11)$$

which should asymptotically coincide with the square of the ground state energy

$$\lim_{n \rightarrow \infty} D_n = e_0^2 = \left(\frac{1}{4} - \zeta_a(1) \right)^2 = 0.1963794236382287 \dots \quad (3.12)$$

From our results of all the density matrix elements for $n = 6$, we can exactly evaluate the dimer-dimer correlation functions D_n for up to $n = 5$:

$$\begin{aligned} D_3 - e_0^2 &= \frac{5}{6}\zeta_a(3) - \frac{4}{3}\zeta_a(1)\zeta_a(3) - \zeta_a(3)^2 - \frac{5}{6}\zeta_a(5) + \frac{10}{3}\zeta_a(1)\zeta_a(5) - \zeta_a(1)^2 \\ &= 0.06082478294036410 \dots, \end{aligned} \quad (3.13)$$

$$\begin{aligned} D_4 - e_0^2 &= \frac{5}{6}\zeta_a(3) - \frac{14}{3}\zeta_a(1)\zeta_a(3) - \frac{43}{3}\zeta_a(3)^2 - \frac{15}{4}\zeta_a(5) + \frac{160}{3}\zeta_a(1)\zeta_a(5) - 5\zeta_a(3)\zeta_a(5) \\ &\quad - \frac{50}{3}\zeta_a(5)^2 + \frac{35}{12}\zeta_a(7) - \frac{140}{3}\zeta_a(1)\zeta_a(7) + 35\zeta_a(3)\zeta_a(7) - \zeta_a(1)^2 \\ &= -0.02773785800119889 \dots, \end{aligned} \quad (3.14)$$

$$\begin{aligned} D_5 - e_0^2 &= \frac{5}{6}\zeta_a(3) - 10\zeta_a(1)\zeta_a(3) - \frac{221}{3}\zeta_a(3)^2 - \frac{320}{9}\zeta_a(3)^2 - \frac{263}{30}\zeta_a(5) + \frac{862}{3}\zeta_a(1)\zeta_a(5) \\ &\quad - \frac{196}{5}\zeta_a(3)\zeta_a(5) + \frac{640}{3}\zeta_a(1)\zeta_a(3)\zeta_a(5) - \frac{904}{15}\zeta_a(3)^2\zeta_a(5) - 1381\zeta_a(5)^2 \\ &\quad + \frac{3616}{3}\zeta_a(1)\zeta_a(5)^2 + \frac{680}{3}\zeta_a(3)\zeta_a(5)^2 + \frac{1040}{9}\zeta_a(5)^3 + \frac{161}{6}\zeta_a(7) - \frac{3220}{3}\zeta_a(1)\zeta_a(7) \\ &\quad + 3038\zeta_a(3)\zeta_a(7) - \frac{6328}{3}\zeta_a(1)\zeta_a(3)\zeta_a(7) - \frac{1904}{3}\zeta_a(3)^2\zeta_a(7) + \frac{1204}{3}\zeta_a(5)\zeta_a(7) \\ &\quad - \frac{1456}{3}\zeta_a(3)\zeta_a(5)\zeta_a(7) - \frac{3773}{3}\zeta_a(7)^2 + \frac{5096}{3}\zeta_a(1)\zeta_a(7)^2 - \frac{189}{10}\zeta_a(9) + 798\zeta_a(1)\zeta_a(9) \\ &\quad - \frac{14224}{5}\zeta_a(3)\zeta_a(9) + 1904\zeta_a(1)\zeta_a(3)\zeta_a(9) + \frac{4368}{5}\zeta_a(3)^2\zeta_a(9) + 2156\zeta_a(5)\zeta_a(9) \\ &\quad - 2912\zeta_a(1)\zeta_a(5)\zeta_a(9) - \zeta_a(1)^2 \\ &= 0.01892813120084483 \dots, \end{aligned} \quad (3.15)$$

The difference $D_n - e_0^2$ also should decay algebraically.

3.4 $AP(n)$: a probability to find an antiferromagnetic string of length n

Emptiness Formation Probability $P(n)$ (1.9) represents the probability to find a ferromagnetic string of length n . Similarly we may consider a probability to find an antiferromagnetic string of length n , which we denote by $AP(n)$

$$AP(n) \equiv P_{+-+ \dots}^{+-+- \dots} + P_{-+-+ \dots}^{--++ \dots}. \quad (3.16)$$

From our results we can exactly evaluate the $AP(n)$ for up to $n = 6$

$$AP(2) = P_{+-}^{+-} + P_{-+}^{-+} = \frac{1}{3} + \frac{2}{3}\zeta_a(1) = 0.7954314537066302 \dots, \quad (3.17)$$

$$AP(3) = P_{+-+}^{+-+-} + P_{-+-}^{-++-} = \frac{1}{6} - \frac{2}{3}\zeta_a(1) + \zeta_a(3) = 0.6061112236630655 \dots, \quad (3.18)$$

$$\begin{aligned}
AP(4) &= P_{+-+-}^{+++-} + P_{-+-+}^{--++} \\
&= \frac{1}{15} - \frac{8}{15}\zeta_a(3) + \frac{4}{3}\zeta_a(1)\zeta_a(3) + \frac{2}{5}\zeta_a(3)^2 + \frac{2}{3}\zeta_a(5) - \frac{4}{3}\zeta_a(1)\zeta_a(5) \\
&= 0.4938083479102196 \dots,
\end{aligned} \tag{3.19}$$

$$\begin{aligned}
AP(5) &= P_{++++}^{+---} + P_{-+++}^{--++} \\
&= \frac{1}{30} - \frac{2}{3}\zeta_a(1) + \frac{91}{15}\zeta_a(3) - \frac{124}{9}\zeta_a(1)\zeta_a(3) - \frac{134}{5}\zeta_a(3)^2 - \frac{355}{18}\zeta_a(5) + \frac{328}{3}\zeta_a(1)\zeta_a(5) \\
&\quad - \frac{38}{9}\zeta_a(3)\zeta_a(5) - \frac{250}{9}\zeta_a(5)^2 + \frac{259}{18}\zeta_a(7) - \frac{854}{9}\zeta_a(1)\zeta_a(7) + \frac{175}{3}\zeta_a(3)\zeta_a(7) \\
&= 0.3943245947356898 \dots,
\end{aligned} \tag{3.20}$$

$$\begin{aligned}
AP(6) &= P_{++++}^{+---} + P_{-+++}^{--++} \\
&= \frac{1}{70} - \frac{1}{5}\zeta_a(1) + \frac{44}{45}\zeta_a(3) + \frac{36}{5}\zeta_a(1)\zeta_a(3) + \frac{2566}{45}\zeta_a(3)^2 \\
&\quad + \frac{4072}{135}\zeta_a(3)^3 + \frac{9754}{1575}\zeta_a(5) - \frac{10984}{45}\zeta_a(1)\zeta_a(5) - \frac{122}{75}\zeta_a(3)\zeta_a(5) \\
&\quad - \frac{8144}{45}\zeta_a(1)\zeta_a(3)\zeta_a(5) + \frac{1368}{25}\zeta_a(3)^2\zeta_a(5) + \frac{374128}{315}\zeta_a(5)^2 - \frac{5472}{5}\zeta_a(1)\zeta_a(5)^2 \\
&\quad - \frac{1868}{9}\zeta_a(3)\zeta_a(5)^2 - \frac{19868}{189}\zeta_a(5)^3 - \frac{1936}{45}\zeta_a(7) + \frac{50848}{45}\zeta_a(1)\zeta_a(7) \\
&\quad - \frac{39794}{15}\zeta_a(3)\zeta_a(7) + \frac{9576}{5}\zeta_a(1)\zeta_a(3)\zeta_a(7) + \frac{26152}{45}\zeta_a(3)^2\zeta_a(7) - \frac{1868}{5}\zeta_a(5)\zeta_a(7) \\
&\quad + \frac{19868}{45}\zeta_a(3)\zeta_a(5)\zeta_a(7) + \frac{47299}{45}\zeta_a(7)^2 - \frac{69538}{45}\zeta_a(1)\zeta_a(7)^2 + \frac{902}{25}\zeta_a(9) \\
&\quad - \frac{4464}{5}\zeta_a(1)\zeta_a(9) + \frac{190052}{75}\zeta_a(3)\zeta_a(9) - \frac{26152}{15}\zeta_a(1)\zeta_a(3)\zeta_a(9) \\
&\quad - \frac{19868}{25}\zeta_a(3)^2\zeta_a(9) - \frac{27028}{15}\zeta_a(5)\zeta_a(9) + \frac{39736}{15}\zeta_a(1)\zeta_a(5)\zeta_a(9) \\
&= 0.3239037769698205 \dots.
\end{aligned} \tag{3.21}$$

It is known that the EFP $P(n)$ shows a Gaussian decay [40, 41]

$$P(n) \simeq An^{-\gamma}C^{-n^2}. \tag{3.22}$$

On the other hand, the analytical asymptotic behavior of $AP(n)$ has not been studied. However, Y. Nishiyama noticed that it decays roughly $AP(n) \simeq [AP(2)]^{n-1}$ from his numerical data before [42]. We compare the formula with our exact data in Table 1.

4 Reduced density matrix and entanglement entropy

At zero temperature, the density matrix for the infinite system can be written as

$$\rho_T \equiv |\text{GS}\rangle\langle\text{GS}|, \tag{4.1}$$

where $|\text{GS}\rangle$ denotes the antiferromagnetic ground state. Let us consider a finite sub-chain of length n in the infinite chain, the rest of which can be regarded as the environment (Figure 1). The reduced density matrix for this sub-chain is obtained by tracing out the environment

Table 1: Asymptotic behavior of $AP(n)$

n	$AP(n)$ Exact	$[AP(2)]^{n-1}$
2	0.795431	0.795431
3	0.606111	0.632711
4	0.493808	0.503278
5	0.394325	0.400323
6	0.323904	0.318430




Figure 1: Finite sub-chain of length n in the infinite chain

from the infinite chain:

$$\rho_n \equiv \text{tr}_E \rho_T = \left[P_{\epsilon_1, \dots, \epsilon_n}^{\epsilon'_1, \dots, \epsilon'_n} \right]_{\epsilon_j, \epsilon'_j = \pm} . \quad (4.2)$$

From our results, we have computed all the eigenvalues ω_α ($\alpha = 1, 2, \dots, 2^n$) of the reduced density matrix ρ_n for up to $n = 6$, which are shown in Figure 2.

We have found that the smallest eigenvalue ω_{2^n} is, for any n , the emptiness formation probability $P(n)$. This is $(n + 1)$ -fold degenerate since the reduced density matrix has the block-diagonalized form by the magnetization of the sub-chain $M_z = \sum_{j=1}^n \epsilon'_j = \sum_{j=1}^n \epsilon_j$, each of which has the non-degenerate smallest eigenvalue $P(n)$. Or in other words, it is a consequence of $SU(2)$ invariance of the Hamiltonian.

From these results we can calculate the von Neumann entropy (Entanglement entropy).

$$S(n) \equiv -\text{tr} \rho_n \log_2 \rho_n = - \sum_{\alpha=1}^{2^n} \omega_\alpha \log_2 \omega_\alpha, \quad (4.3)$$

which is considered to measure how the ground state is entangled [43]. The exact numerical values of $S(n)$ up to $n = 6$ are shown in Table 2. It can be seen that as n grows the von Neumann entropy is increasing in small steps. The asymptotic behavior of this entropy is discussed in [43]. In the massive region $\Delta > 1$, the von Neumann entropy will be saturated as n grows, which means the ground state is well approximated by a subsystem of a finite length corresponding to the large eigenvalues of reduced density matrix. On the other hand, in the massless case $-1 < \Delta \leq 1$, the conformal field theory predict that the von Neumann entropy shows a logarithmic divergence [44], of which the explicit form for the XXX case $\Delta = 1$ reads

$$S(n) \simeq \frac{1}{3} \log_2 n + k. \quad (4.4)$$




Figure 2: Eigenvalue-distribution of density matrices

Table 2: von Neumann entropy $S(n)$ of a finite sub-chain of length n			
$S(1)$	$S(2)$	$S(3)$	$S(4)$
1	1.3758573262887466	1.5824933209573855	1.7247050949099274

$S(5)$	$S(6)$
1.833704916848315	1.922358833819333

We estimate the numerical value of the constant term $k = 1.06209$ by the extrapolation $S(n) - \frac{1}{3} \log_2 n = c_0 + c_1/n + c_2/n^2$. Our exact results for up to $n = 6$ agree quite well with the asymptotic formula as shown in Figure 3.

5 Summary and discussion

We have succeeded in obtaining the analytical forms of all the density matrix elements on six lattice sites for Heisenberg chain using the algebraic method based on the qKZ equations developed in [34]. We have obtained several physically interesting correlation functions, such as chiral correlation functions, dimer-dimer correlation functions, etc. . . . Subsequently we have calculated all the eigenvalues of the reduced density matrix ρ_n for up to $n = 6$. We observe that the smallest eigenvalue is the emptiness formation probability $P(n)$ and is $(n + 1)$ -fold degenerate. Of course, it may be more desirable if we could describe the eigenvector corresponding to the largest eigenvalue. From these results we have computed the von Neumann entropy, which shows a good agreement with the asymptotic formula derived via the conformal field theory.

Acknowledgement

The authors are grateful to H. Boos, V.E. Korepin, Y. Nishiyama and K. Sakai for fruitful collaborations of early stage of this work. Especially we thank V.E. Korepin for suggesting us to calculate the entanglement entropy. We also thank M. Bortz and M. Batchelor for valuable discussions. This work is in part supported by Grant-in-Aid for the Scientific Research (B) No. 18340112. from the Ministry of Education, Culture, Sports, Science and Technology, Japan. It is also supported by JSPS and ARC under the Japan-Australian joint project in cooperative science program.

Appendix A Known results for up to $n = 5$

In this appendix we present all the independent correlation functions for up to $n = 5$ except for the two-point correlation functions $\langle S_1^z S_n^z \rangle$ given in Introduction. First we give the correlation functions on four lattice sites [19]:

$$\begin{aligned}
\langle S_1^x S_2^x S_3^z S_4^z \rangle &= \frac{1}{240} + \frac{1}{12} \zeta_a(1) - \frac{91}{180} \zeta_a(3) + \frac{2}{9} \zeta_a(1) \zeta_a(3) + \frac{1}{15} \zeta_a(3)^2 + \frac{7}{18} \zeta_a(5) - \frac{2}{9} \zeta_a(1) \zeta_a(5) \\
&= 0.02750969925180030 \dots,
\end{aligned} \tag{A.5}$$




Figure 3: von Neumann entropy $S(n)$ of a finite sub-chain of length n

$$\begin{aligned} \langle S_1^x S_2^z S_3^x S_4^z \rangle &= \frac{1}{240} - \frac{1}{6}\zeta_a(1) + \frac{77}{90}\zeta_a(3) - \frac{5}{9}\zeta_a(1)\zeta_a(3) - \frac{4}{15}\zeta_a(3)^2 - \frac{13}{18}\zeta_a(5) + \frac{8}{9}\zeta_a(1)\zeta_a(5) \\ &= -0.007080326815007911 \dots, \end{aligned} \quad (\text{A.6})$$

$$\begin{aligned} \langle S_1^x S_2^z S_3^z S_4^x \rangle &= \frac{1}{240} - \frac{1}{4}\zeta_a(1) + \frac{169}{180}\zeta_a(3) - \frac{5}{9}\zeta_a(1)\zeta_a(3) - \frac{4}{15}\zeta_a(3)^2 - \frac{13}{18}\zeta_a(5) + \frac{8}{9}\zeta_a(1)\zeta_a(5) \\ &= 0.01028596458580462 \dots, \end{aligned} \quad (\text{A.7})$$

$$\langle S_1^z S_2^z S_3^z S_4^z \rangle = \langle S_1^x S_2^z S_3^z S_4^x \rangle + \langle S_1^x S_2^z S_3^x S_4^z \rangle + \langle S_1^x S_2^z S_3^z S_4^x \rangle. \quad (\text{A.8})$$

Note that on the antiferromagnetic ground state without magnetic field, the correlation functions with an odd number of spin operators vanish. Considering the isotropy of the Hamiltonian (1.1), we can see that the independent correlation functions on four lattice sites are restricted to the above ones and the two-point correlation function $\langle S_1^z S_4^z \rangle$.

Correspondingly the independent correlation functions on five lattice sites are [34]

$$\begin{aligned} \langle S_1^x S_2^x S_3^z S_5^z \rangle &= \frac{1}{240} + \frac{1}{12}\zeta_a(1) - \frac{517}{360}\zeta_a(3) + \frac{25}{9}\zeta_a(1)\zeta_a(3) + \frac{203}{45}\zeta_a(3)^2 + \frac{305}{72}\zeta_a(5) \\ &\quad - \frac{172}{9}\zeta_a(1)\zeta_a(5) + \frac{4}{9}\zeta_a(3)\zeta_a(5) + \frac{40}{9}\zeta_a(5)^2 - \frac{35}{12}\zeta_a(7) + \frac{49}{3}\zeta_a(1)\zeta_a(7) \\ &\quad - \frac{28}{3}\zeta_a(3)\zeta_a(7) = -0.009892435084700294 \dots, \end{aligned} \quad (\text{A.9})$$

$$\begin{aligned}
\langle S_1^x S_2^z S_3^x S_5^z \rangle &= \frac{1}{240} - \frac{1}{4}\zeta_a(1) + \frac{301}{120}\zeta_a(3) - 6\zeta_a(1)\zeta_a(3) - \frac{1079}{90}\zeta_a(3)^2 - \frac{65}{8}\zeta_a(5) \\
&\quad + \frac{146}{3}\zeta_a(1)\zeta_a(5) - \frac{37}{18}\zeta_a(3)\zeta_a(5) - \frac{110}{9}\zeta_a(5)^2 + \frac{35}{6}\zeta_a(7) - 42\zeta_a(1)\zeta_a(7) \\
&\quad + \frac{77}{3}\zeta_a(3)\zeta_a(7) = 0.002788973399542967 \dots, \tag{A.10}
\end{aligned}$$

$$\begin{aligned}
\langle S_1^x S_2^z S_3^z S_5^x \rangle &= \frac{1}{240} - \frac{5}{12}\zeta_a(1) + \frac{569}{180}\zeta_a(3) - \frac{61}{9}\zeta_a(1)\zeta_a(3) - \frac{1109}{90}\zeta_a(3)^2 - \frac{155}{18}\zeta_a(5) \\
&\quad + \frac{448}{9}\zeta_a(1)\zeta_a(5) - \frac{37}{18}\zeta_a(3)\zeta_a(5) - \frac{110}{9}\zeta_a(5)^2 + \frac{35}{6}\zeta_a(7) - 42\zeta_a(1)\zeta_a(7) \\
&\quad + \frac{77}{3}\zeta_a(3)\zeta_a(7) = -0.005056660894819286 \dots, \tag{A.11}
\end{aligned}$$

$$\langle S_1^z S_2^z S_3^z S_5^z \rangle = \langle S_1^x S_2^x S_3^z S_5^z \rangle + \langle S_1^x S_2^z S_3^x S_5^z \rangle + \langle S_1^x S_2^z S_3^z S_5^x \rangle, \tag{A.12}$$

$$\begin{aligned}
\langle S_1^x S_2^x S_4^z S_5^z \rangle &= \frac{1}{240} + \frac{1}{4}\zeta_a(1) - \frac{419}{120}\zeta_a(3) + \frac{70}{9}\zeta_a(1)\zeta_a(3) + \frac{221}{15}\zeta_a(3)^2 + \frac{889}{72}\zeta_a(5) \\
&\quad - \frac{184}{3}\zeta_a(1)\zeta_a(5) + \frac{17}{9}\zeta_a(3)\zeta_a(5) + \frac{130}{9}\zeta_a(5)^2 - \frac{329}{36}\zeta_a(7) + \frac{476}{9}\zeta_a(1)\zeta_a(7) \\
&\quad - \frac{91}{3}\zeta_a(3)\zeta_a(7) = 0.01857662093837095 \dots, \tag{A.13}
\end{aligned}$$

$$\begin{aligned}
\langle S_1^x S_2^z S_4^x S_5^z \rangle &= \frac{1}{240} - \frac{5}{12}\zeta_a(1) + \frac{1883}{360}\zeta_a(3) - \frac{112}{9}\zeta_a(1)\zeta_a(3) - \frac{1102}{45}\zeta_a(3)^2 - \frac{457}{24}\zeta_a(5) \\
&\quad + \frac{908}{9}\zeta_a(1)\zeta_a(5) - \frac{11}{3}\zeta_a(3)\zeta_a(5) - \frac{220}{9}\zeta_a(5)^2 + \frac{511}{36}\zeta_a(7) - \frac{784}{9}\zeta_a(1)\zeta_a(7) \\
&\quad + \frac{154}{3}\zeta_a(3)\zeta_a(7) = -0.001089368972914637 \dots, \tag{A.14}
\end{aligned}$$

$$\begin{aligned}
\langle S_1^x S_2^z S_4^z S_5^x \rangle &= \frac{1}{240} - \frac{1}{2}\zeta_a(1) + \frac{497}{90}\zeta_a(3) - \frac{112}{9}\zeta_a(1)\zeta_a(3) - \frac{1102}{45}\zeta_a(3)^2 - \frac{77}{4}\zeta_a(5) \\
&\quad + \frac{908}{9}\zeta_a(1)\zeta_a(5) - \frac{11}{3}\zeta_a(3)\zeta_a(5) - \frac{220}{9}\zeta_a(5)^2 + \frac{511}{36}\zeta_a(7) - \frac{784}{9}\zeta_a(1)\zeta_a(7) \\
&\quad + \frac{154}{3}\zeta_a(3)\zeta_a(7) = 0.001573361370145065 \dots, \tag{A.15}
\end{aligned}$$

$$\langle S_1^z S_2^z S_4^z S_5^z \rangle = \langle S_1^x S_2^x S_4^z S_5^z \rangle + \langle S_1^x S_2^z S_4^x S_5^z \rangle + \langle S_1^x S_2^z S_4^z S_5^x \rangle. \tag{A.16}$$

Appendix B Explicit form for all the correlation functions on 6 lattice sites

In this appendix the explicit results for all the correlation functions on 6 lattice sites are shown. There are 24 independent correlation functions among them. Any other correlation functions can be computed by the following relations:

$$\langle S_1^z S_3^z S_4^z S_6^z \rangle = \langle S_1^x S_3^x S_4^z S_6^z \rangle + \langle S_1^x S_3^z S_4^x S_6^z \rangle + \langle S_1^x S_3^z S_4^z S_6^x \rangle, \tag{B.1}$$

$$\langle S_1^z S_2^z S_5^z S_6^z \rangle = \langle S_1^x S_2^x S_5^z S_6^z \rangle + \langle S_1^x S_2^z S_5^x S_6^z \rangle + \langle S_1^x S_2^z S_5^z S_6^x \rangle, \tag{B.2}$$

$$\langle S_1^z S_4^z S_5^z S_6^z \rangle = \langle S_1^x S_4^x S_5^z S_6^z \rangle + \langle S_1^x S_4^z S_5^x S_6^z \rangle + \langle S_1^x S_4^z S_5^z S_6^x \rangle, \tag{B.3}$$

$$\langle S_1^z S_3^z S_5^z S_6^z \rangle = \langle S_1^x S_3^x S_5^z S_6^z \rangle + \langle S_1^x S_3^z S_5^x S_6^z \rangle + \langle S_1^x S_3^z S_5^z S_6^x \rangle, \tag{B.4}$$

$$\langle S_1^x S_2^x S_3^y S_4^y S_5^z S_6^z \rangle = \langle S_1^x S_2^x S_3^x S_4^x S_5^y S_6^y \rangle - \langle S_1^x S_2^x S_3^y S_4^y S_5^y S_6^z \rangle - \langle S_1^x S_2^x S_3^y S_4^z S_5^z S_6^y \rangle, \quad (\text{B.5})$$

$$\langle S_1^x S_2^y S_3^x S_4^z S_5^y S_6^z \rangle = \langle S_1^x S_2^x S_3^x S_4^x S_5^y S_6^y \rangle - \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle - \langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle, \quad (\text{B.6})$$

$$\langle S_1^x S_2^y S_3^y S_4^z S_5^y S_6^z \rangle = \langle S_1^x S_2^x S_3^x S_4^x S_5^y S_6^y \rangle - \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle - \langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle, \quad (\text{B.7})$$

$$\langle S_1^x S_2^y S_3^z S_4^y S_5^y S_6^z \rangle = \langle S_1^x S_2^x S_3^x S_4^x S_5^y S_6^y \rangle - \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle - \langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle, \quad (\text{B.8})$$

$$\langle S_1^x S_2^y S_3^z S_4^z S_5^y S_6^x \rangle = \langle S_1^x S_2^x S_3^x S_4^x S_5^y S_6^y \rangle - \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle - \langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle, \quad (\text{B.9})$$

$$\langle S_1^x S_2^y S_3^z S_4^z S_5^x S_6^y \rangle = \langle S_1^x S_2^x S_3^x S_4^x S_5^y S_6^y \rangle - \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle - \langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle, \quad (\text{B.10})$$

$$\begin{aligned} \langle S_1^x S_2^x S_3^y S_4^y S_5^x S_6^x \rangle &= \langle S_1^x S_2^y S_3^z S_4^z S_5^y S_6^x \rangle + \langle S_1^x S_2^x S_3^x S_4^x S_5^y S_6^y \rangle + \langle S_1^x S_2^y S_3^x S_4^x S_5^x S_6^y \rangle \\ &- \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle - \langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle - \langle S_1^x S_2^y S_3^z S_4^z S_5^z S_6^y \rangle, \end{aligned} \quad (\text{B.11})$$

$$\begin{aligned} \langle S_1^x S_2^y S_3^x S_4^y S_5^x S_6^x \rangle &= \langle S_1^x S_2^y S_3^z S_4^z S_5^y S_6^x \rangle + \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle + \langle S_1^x S_2^y S_3^x S_4^x S_5^x S_6^y \rangle \\ &- \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle - \langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle - \langle S_1^x S_2^y S_3^z S_4^z S_5^z S_6^y \rangle, \end{aligned} \quad (\text{B.12})$$

$$\begin{aligned} \langle S_1^x S_2^y S_3^y S_4^y S_5^x S_6^x \rangle &= \langle S_1^x S_2^y S_3^z S_4^z S_5^y S_6^x \rangle + \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle + \langle S_1^x S_2^y S_3^x S_4^x S_5^x S_6^y \rangle \\ &- \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle - \langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle - \langle S_1^x S_2^y S_3^z S_4^z S_5^z S_6^y \rangle, \end{aligned} \quad (\text{B.13})$$

$$\begin{aligned} \langle S_1^x S_2^x S_3^x S_4^x S_5^x S_6^x \rangle &= \langle S_1^x S_2^y S_3^z S_4^z S_5^y S_6^x \rangle + \langle S_1^x S_2^x S_3^x S_4^x S_5^y S_6^y \rangle + \langle S_1^x S_2^y S_3^x S_4^x S_5^x S_6^y \rangle \\ &+ \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle + \langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle + \langle S_1^x S_2^y S_3^z S_4^z S_5^z S_6^y \rangle \\ &- \langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle - \langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle - \langle S_1^x S_2^y S_3^z S_4^z S_5^z S_6^y \rangle. \end{aligned} \quad (\text{B.14})$$

Below we shall give the explicit results for the independent ones.

$$\begin{aligned} \langle S_1^z S_6^z \rangle &= \frac{1}{12} - \frac{25}{3} \zeta_a(1) + \frac{800}{9} \zeta_a(3) - \frac{1192}{3} \zeta_a(1) \zeta_a(3) - \frac{15368}{9} \zeta_a(3)^2 \\ &- 608 \zeta_a(3)^3 - \frac{4228}{9} \zeta_a(5) + \frac{64256}{9} \zeta_a(1) \zeta_a(5) - \frac{976}{9} \zeta_a(3) \zeta_a(5) \\ &+ 3648 \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{3328}{3} \zeta_a(3)^2 \zeta_a(5) - \frac{76640}{3} \zeta_a(5)^2 \\ &+ \frac{66560}{3} \zeta_a(1) \zeta_a(5)^2 + \frac{12640}{3} \zeta_a(3) \zeta_a(5)^2 + \frac{6400}{3} \zeta_a(5)^3 + \frac{9674}{9} \zeta_a(7) \\ &+ 56952 \zeta_a(3) \zeta_a(7) - \frac{225848}{9} \zeta_a(1) \zeta_a(7) - \frac{116480}{3} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\ &- \frac{35392}{3} \zeta_a(3)^2 \zeta_a(7) + 7840 \zeta_a(5) \zeta_a(7) - 8960 \zeta_a(3) \zeta_a(5) \zeta_a(7) \\ &- \frac{66640}{3} \zeta_a(7)^2 + 31360 \zeta_a(1) \zeta_a(7)^2 - 686 \zeta_a(9) \\ &+ 18368 \zeta_a(1) \zeta_a(9) - 53312 \zeta_a(3) \zeta_a(9) + 35392 \zeta_a(1) \zeta_a(3) \zeta_a(9) \\ &+ 16128 \zeta_a(3)^2 \zeta_a(9) + 38080 \zeta_a(5) \zeta_a(9) - 53760 \zeta_a(1) \zeta_a(5) \zeta_a(9) \\ &= -0.03089036664760932 \dots \end{aligned} \quad (\text{B.15})$$

$$\begin{aligned}
\langle S_1^x S_3^x S_4^z S_6^z \rangle &= \frac{1}{240} + \frac{1}{10} \zeta_a(1) - \frac{259}{90} \zeta_a(3) + \frac{463}{45} \zeta_a(1) \zeta_a(3) + \frac{1951}{45} \zeta_a(3)^2 \\
&+ \frac{2404}{135} \zeta_a(3)^3 + \frac{1348}{75} \zeta_a(5) - \frac{8918}{45} \zeta_a(1) \zeta_a(5) - \frac{3127}{225} \zeta_a(3) \zeta_a(5) \\
&- \frac{4808}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) + \frac{7804}{225} \zeta_a(3)^2 \zeta_a(5) + \frac{33598}{45} \zeta_a(5)^2 \\
&- \frac{31216}{45} \zeta_a(1) \zeta_a(5)^2 - \frac{1196}{9} \zeta_a(3) \zeta_a(5)^2 - \frac{1808}{27} \zeta_a(5)^3 - \frac{413}{9} \zeta_a(7) \\
&+ \frac{36421}{45} \zeta_a(1) \zeta_a(7) - \frac{8393}{5} \zeta_a(3) \zeta_a(7) + \frac{54628}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&+ \frac{16744}{45} \zeta_a(3)^2 \zeta_a(7) - \frac{10864}{45} \zeta_a(5) \zeta_a(7) + \frac{12656}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&+ \frac{29008}{45} \zeta_a(7)^2 - \frac{44296}{45} \zeta_a(1) \zeta_a(7)^2 + \frac{1533}{50} \zeta_a(9) \\
&- \frac{3108}{5} \zeta_a(1) \zeta_a(9) + \frac{120344}{75} \zeta_a(3) \zeta_a(9) - \frac{16744}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&- \frac{12656}{25} \zeta_a(3)^2 \zeta_a(9) - \frac{16576}{15} \zeta_a(5) \zeta_a(9) + \frac{25312}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= 0.003681507672875026 \dots
\end{aligned} \tag{B.16}$$

$$\begin{aligned}
\langle S_1^x S_3^z S_4^x S_6^z \rangle &= \frac{1}{240} - \frac{19}{60} \zeta_a(1) + \frac{1007}{180} \zeta_a(3) - \frac{1252}{45} \zeta_a(1) \zeta_a(3) - \frac{1978}{15} \zeta_a(3)^2 \\
&- \frac{6256}{135} \zeta_a(3)^3 - \frac{979}{25} \zeta_a(5) + \frac{25322}{45} \zeta_a(1) \zeta_a(5) + \frac{538}{225} \zeta_a(3) \zeta_a(5) \\
&+ \frac{12512}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{6292}{75} \zeta_a(3)^2 \zeta_a(5) - \frac{85687}{45} \zeta_a(5)^2 \\
&+ \frac{25168}{15} \zeta_a(1) \zeta_a(5)^2 + \frac{2864}{9} \zeta_a(3) \zeta_a(5)^2 + \frac{4352}{27} \zeta_a(5)^3 + \frac{1771}{18} \zeta_a(7) \\
&- \frac{89929}{45} \zeta_a(1) \zeta_a(7) + \frac{191513}{45} \zeta_a(3) \zeta_a(7) - \frac{44044}{15} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&- \frac{40096}{45} \zeta_a(3)^2 \zeta_a(7) + \frac{26866}{45} \zeta_a(5) \zeta_a(7) - \frac{30464}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&- \frac{73402}{45} \zeta_a(7)^2 + \frac{106624}{45} \zeta_a(1) \zeta_a(7)^2 - \frac{3227}{50} \zeta_a(9) \\
&+ \frac{7322}{5} \zeta_a(1) \zeta_a(9) - \frac{298886}{75} \zeta_a(3) \zeta_a(9) + \frac{40096}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&+ \frac{30464}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{41944}{15} \zeta_a(5) \zeta_a(9) - \frac{60928}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= -0.001116347734065082 \dots
\end{aligned} \tag{B.17}$$

$$\begin{aligned}
\langle S_1^x S_3^z S_4^z S_6^x \rangle &= \frac{1}{240} - \frac{13}{20} \zeta_a(1) + \frac{479}{60} \zeta_a(3) - \frac{1537}{45} \zeta_a(1) \zeta_a(3) - \frac{6709}{45} \zeta_a(3)^2 \\
&\quad - \frac{2452}{45} \zeta_a(3)^3 - \frac{10201}{225} \zeta_a(5) + \frac{9524}{15} \zeta_a(1) \zeta_a(5) + \frac{37}{25} \zeta_a(3) \zeta_a(5) \\
&\quad + \frac{4904}{15} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{22616}{225} \zeta_a(3)^2 \zeta_a(5) - \frac{102592}{45} \zeta_a(5)^2 \\
&\quad + \frac{90464}{45} \zeta_a(1) \zeta_a(5)^2 + \frac{1148}{3} \zeta_a(3) \zeta_a(5)^2 + \frac{1744}{9} \zeta_a(5)^3 + \frac{973}{9} \zeta_a(7) \\
&\quad - \frac{103544}{45} \zeta_a(1) \zeta_a(7) + \frac{229138}{45} \zeta_a(3) \zeta_a(7) - \frac{158312}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&\quad - \frac{16072}{15} \zeta_a(3)^2 \zeta_a(7) + \frac{31906}{45} \zeta_a(5) \zeta_a(7) - \frac{12208}{15} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&\quad - \frac{9898}{5} \zeta_a(7)^2 + \frac{42728}{15} \zeta_a(1) \zeta_a(7)^2 - \frac{3507}{50} \zeta_a(9) \\
&\quad + \frac{8512}{5} \zeta_a(1) \zeta_a(9) - \frac{119742}{25} \zeta_a(3) \zeta_a(9) + \frac{16072}{5} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&\quad + \frac{36624}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{16968}{5} \zeta_a(5) \zeta_a(9) - \frac{24416}{5} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= 0.003069653070471227 \dots
\end{aligned} \tag{B.18}$$

$$\begin{aligned}
\langle S_1^x S_2^x S_5^z S_6^z \rangle &= \frac{1}{240} + \frac{29}{60} \zeta_a(1) - \frac{568}{45} \zeta_a(3) + \frac{1024}{15} \zeta_a(1) \zeta_a(3) + \frac{15863}{45} \zeta_a(3)^2 \\
&\quad + \frac{5872}{45} \zeta_a(3)^3 + \frac{25486}{225} \zeta_a(5) - \frac{69922}{45} \zeta_a(1) \zeta_a(5) - \frac{13598}{225} \zeta_a(3) \zeta_a(5) \\
&\quad - \frac{11744}{15} \zeta_a(1) \zeta_a(3) \zeta_a(5) + \frac{53816}{225} \zeta_a(3)^2 \zeta_a(5) + \frac{246547}{45} \zeta_a(5)^2 \\
&\quad - \frac{215264}{45} \zeta_a(1) \zeta_a(5)^2 - \frac{2728}{3} \zeta_a(3) \zeta_a(5)^2 - \frac{4144}{9} \zeta_a(5)^3 - \frac{61327}{180} \zeta_a(7) \\
&\quad + \frac{271264}{45} \zeta_a(1) \zeta_a(7) - \frac{61474}{5} \zeta_a(3) \zeta_a(7) + \frac{376712}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&\quad + \frac{38192}{15} \zeta_a(3)^2 \zeta_a(7) - \frac{79436}{45} \zeta_a(5) \zeta_a(7) + \frac{29008}{15} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&\quad + \frac{23373}{5} \zeta_a(7)^2 - \frac{101528}{15} \zeta_a(1) \zeta_a(7)^2 + \frac{11977}{50} \zeta_a(9) \\
&\quad - \frac{22722}{5} \zeta_a(1) \zeta_a(9) + \frac{290752}{25} \zeta_a(3) \zeta_a(9) - \frac{38192}{5} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&\quad - \frac{87024}{25} \zeta_a(3)^2 \zeta_a(9) - \frac{40068}{5} \zeta_a(5) \zeta_a(9) + \frac{58016}{5} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= 0.02384723373803033 \dots
\end{aligned} \tag{B.19}$$

$$\begin{aligned}
\langle S_1^x S_2^z S_5^x S_6^z \rangle &= \frac{1}{240} - \frac{23}{30} \zeta_a(1) + \frac{563}{30} za3 - \frac{4678}{45} \zeta_a(1) \zeta_a(3) - \frac{8114}{15} \zeta_a(3)^2 \\
&- \frac{27224}{135} \zeta_a(3)^3 - \frac{38389}{225} \zeta_a(5) + \frac{35666}{15} \zeta_a(1) \zeta_a(5) + \frac{18902}{225} \zeta_a(3) \zeta_a(5) \\
&+ \frac{54448}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{82984}{225} \zeta_a(3)^2 \zeta_a(5) - \frac{42242}{5} \zeta_a(5)^2 \\
&+ \frac{331936}{45} \zeta_a(1) \zeta_a(5)^2 + \frac{12616}{9} \zeta_a(3) \zeta_a(5)^2 + \frac{19168}{27} \zeta_a(5)^3 + \frac{23177}{45} \zeta_a(7) \\
&- \frac{414911}{45} \zeta_a(1) \zeta_a(7) + \frac{284228}{15} \zeta_a(3) \zeta_a(7) - \frac{580888}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&- \frac{176624}{45} \zeta_a(3)^2 \zeta_a(7) + \frac{122164}{45} \zeta_a(5) \zeta_a(7) - \frac{134176}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&- \frac{324968}{45} \zeta_a(7)^2 + \frac{469616}{45} \zeta_a(1) \zeta_a(7)^2 - \frac{18123}{50} \zeta_a(9) \\
&+ \frac{34748}{5} \zeta_a(1) \zeta_a(9) - \frac{1343944}{75} \zeta_a(3) \zeta_a(9) + \frac{176624}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&+ \frac{134176}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{185696}{15} \zeta_a(5) \zeta_a(9) - \frac{268352}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= -0.001365096861940014 \dots
\end{aligned} \tag{B.20}$$

$$\begin{aligned}
\langle S_1^x S_2^z S_5^z S_6^x \rangle &= \frac{1}{240} - \frac{17}{20} \zeta_a(1) + \frac{872}{45} \zeta_a(3) - \frac{4688}{45} \zeta_a(1) \zeta_a(3) - \frac{24352}{45} \zeta_a(3)^2 \\
&- \frac{27224}{135} \zeta_a(3)^3 - \frac{77453}{450} \zeta_a(5) + \frac{107078}{45} \zeta_a(1) \zeta_a(5) + \frac{18952}{225} \zeta_a(3) \zeta_a(5) \\
&+ \frac{54448}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{82984}{225} \zeta_a(3)^2 \zeta_a(5) - \frac{42242}{5} \zeta_a(5)^2 \\
&+ \frac{331936}{45} \zeta_a(1) \zeta_a(5)^2 + \frac{12616}{9} \zeta_a(3) \zeta_a(5)^2 + \frac{19168}{27} \zeta_a(5)^3 + \frac{30961}{60} \zeta_a(7) \\
&- \frac{46109}{5} \zeta_a(1) \zeta_a(7) + \frac{284228}{15} \zeta_a(3) \zeta_a(7) - \frac{580888}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&- \frac{176624}{45} \zeta_a(3)^2 \zeta_a(7) + \frac{122164}{45} \zeta_a(5) \zeta_a(7) - \frac{134176}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&- \frac{324968}{45} \zeta_a(7)^2 + \frac{469616}{45} \zeta_a(1) \zeta_a(7)^2 - \frac{18123}{50} \zeta_a(9) \\
&+ \frac{34748}{5} \zeta_a(1) \zeta_a(9) - \frac{1343944}{75} \zeta_a(3) \zeta_a(9) + \frac{176624}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&+ \frac{134176}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{185696}{15} \zeta_a(5) \zeta_a(9) - \frac{268352}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= 0.001592580594206881 \dots
\end{aligned} \tag{B.21}$$

$$\begin{aligned}
\langle S_1^x S_4^x S_5^z S_6^z \rangle &= \frac{1}{240} + \frac{1}{20} \zeta_a(1) - \frac{1093}{360} \zeta_a(3) + \frac{659}{45} \zeta_a(1) \zeta_a(3) + \frac{5621}{90} \zeta_a(3)^2 \\
&+ \frac{2704}{135} \zeta_a(3)^3 + \frac{38411}{1800} \zeta_a(5) - \frac{2452}{9} \zeta_a(1) \zeta_a(5) - \frac{3857}{450} \zeta_a(3) \zeta_a(5) \\
&- \frac{5408}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) + \frac{8272}{225} \zeta_a(3)^2 \zeta_a(5) + \frac{39289}{45} \zeta_a(5)^2 \\
&- \frac{33088}{45} \zeta_a(1) \zeta_a(5)^2 - \frac{1256}{9} \zeta_a(3) \zeta_a(5)^2 - \frac{1904}{27} \zeta_a(5)^3 - \frac{1589}{30} \zeta_a(7) \\
&+ \frac{14581}{15} \zeta_a(1) \zeta_a(7) - \frac{17633}{9} \zeta_a(3) \zeta_a(7) + \frac{57904}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&+ \frac{17584}{45} \zeta_a(3)^2 \zeta_a(7) - \frac{1393}{5} \zeta_a(5) \zeta_a(7) + \frac{13328}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&+ \frac{33124}{45} \zeta_a(7)^2 - \frac{46648}{45} \zeta_a(1) \zeta_a(7)^2 + \frac{1729}{50} \zeta_a(9) \\
&- 714 \zeta_a(1) \zeta_a(9) + \frac{137732}{75} \zeta_a(3) \zeta_a(9) - \frac{17584}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&- \frac{13328}{25} \zeta_a(3)^2 \zeta_a(9) - \frac{18928}{15} \zeta_a(5) \zeta_a(9) + \frac{26656}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= 0.009188091173609528 \dots
\end{aligned} \tag{B.22}$$

$$\begin{aligned}
\langle S_1^x S_4^z S_5^x S_6^z \rangle &= \frac{1}{240} - \frac{11}{30} \zeta_a(1) + \frac{254}{45} \zeta_a(3) - \frac{144}{5} \zeta_a(1) \zeta_a(3) - \frac{6517}{45} \zeta_a(3)^2 \\
&- \frac{7556}{135} \zeta_a(3)^3 - \frac{18091}{450} \zeta_a(5) + \frac{5578}{9} \zeta_a(1) \zeta_a(5) + \frac{253}{75} \zeta_a(3) \zeta_a(5) \\
&+ \frac{15112}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{22928}{225} \zeta_a(3)^2 \zeta_a(5) - \frac{105686}{45} \zeta_a(5)^2 \\
&+ \frac{91712}{45} \zeta_a(1) \zeta_a(5)^2 + \frac{3484}{9} \zeta_a(3) \zeta_a(5)^2 + \frac{5296}{27} \zeta_a(5)^3 + \frac{9793}{90} \zeta_a(7) \\
&- \frac{106547}{45} \zeta_a(1) \zeta_a(7) + \frac{47236}{9} \zeta_a(3) \zeta_a(7) - \frac{160496}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&- \frac{48776}{45} \zeta_a(3)^2 \zeta_a(7) + \frac{3682}{5} \zeta_a(5) \zeta_a(7) - \frac{37072}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&- \frac{91826}{45} \zeta_a(7)^2 + \frac{129752}{45} \zeta_a(1) \zeta_a(7)^2 - \frac{1848}{25} \zeta_a(9) \\
&+ 1778 \zeta_a(1) \zeta_a(9) - \frac{371518}{75} \zeta_a(3) \zeta_a(9) + \frac{48776}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&+ \frac{37072}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{52472}{15} \zeta_a(5) \zeta_a(9) - \frac{74144}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= -0.003158274321296133 \dots
\end{aligned} \tag{B.23}$$

$$\begin{aligned}
\langle S_1^x S_4^z S_5^z S_6^x \rangle = & \frac{1}{240} - \frac{37}{60} \zeta_a(1) + \frac{2887}{360} \zeta_a(3) - \frac{1651}{45} \zeta_a(1) \zeta_a(3) - \frac{4813}{30} \zeta_a(3)^2 \\
& - \frac{7556}{135} \zeta_a(3)^3 - \frac{28313}{600} \zeta_a(5) + \frac{6124}{9} \zeta_a(1) \zeta_a(5) - \frac{7}{450} \zeta_a(3) \zeta_a(5) \\
& + \frac{15112}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{22928}{225} \zeta_a(3)^2 \zeta_a(5) - \frac{106436}{45} \zeta_a(5)^2 \\
& + \frac{91712}{45} \zeta_a(1) \zeta_a(5)^2 + \frac{3484}{9} \zeta_a(3) \zeta_a(5)^2 + \frac{5296}{27} \zeta_a(5)^3 + \frac{20461}{180} \zeta_a(7) \\
& - \frac{108892}{45} \zeta_a(1) \zeta_a(7) + \frac{47551}{9} \zeta_a(3) \zeta_a(7) - \frac{160496}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
& - \frac{48776}{45} \zeta_a(3)^2 \zeta_a(7) + \frac{3682}{5} \zeta_a(5) \zeta_a(7) - \frac{37072}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
& - \frac{91826}{45} \zeta_a(7)^2 + \frac{129752}{45} \zeta_a(1) \zeta_a(7)^2 - \frac{1848}{25} \zeta_a(9) \\
& + 1778 \zeta_a(1) \zeta_a(9) - \frac{371518}{75} \zeta_a(3) \zeta_a(9) + \frac{48776}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
& + \frac{37072}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{52472}{15} \zeta_a(5) \zeta_a(9) - \frac{74144}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
& = 0.005680615367538651 \dots
\end{aligned} \tag{B.24}$$

$$\begin{aligned}
\langle S_1^x S_3^x S_5^z S_6^z \rangle = & \frac{1}{240} + \frac{17}{60} \zeta_a(1) - \frac{2617}{360} \zeta_a(3) + \frac{1546}{45} \zeta_a(1) \zeta_a(3) + \frac{2935}{18} \zeta_a(3)^2 \\
& + \frac{8008}{135} \zeta_a(3)^3 + \frac{10811}{200} \zeta_a(5) - \frac{6430}{9} \zeta_a(1) \zeta_a(5) - \frac{3481}{150} \zeta_a(3) \zeta_a(5) \\
& - \frac{16016}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) + \frac{8216}{75} \zeta_a(3)^2 \zeta_a(5) + \frac{111916}{45} \zeta_a(5)^2 \\
& - \frac{32864}{15} \zeta_a(1) \zeta_a(5)^2 - \frac{3752}{9} \zeta_a(3) \zeta_a(5)^2 - \frac{5696}{27} \zeta_a(5)^3 - \frac{13097}{90} \zeta_a(7) \\
& + \frac{121282}{45} \zeta_a(1) \zeta_a(7) - \frac{16730}{3} \zeta_a(3) \zeta_a(7) + \frac{57512}{15} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
& + \frac{52528}{45} \zeta_a(3)^2 \zeta_a(7) - \frac{11921}{15} \zeta_a(5) \zeta_a(7) + \frac{39872}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
& + \frac{95746}{45} \zeta_a(7)^2 - \frac{139552}{45} \zeta_a(1) \zeta_a(7)^2 + \frac{4921}{50} \zeta_a(9) \\
& - 2016 \zeta_a(1) \zeta_a(9) + \frac{394898}{75} \zeta_a(3) \zeta_a(9) - \frac{52528}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
& - \frac{39872}{25} \zeta_a(3)^2 \zeta_a(9) - \frac{54712}{15} \zeta_a(5) \zeta_a(9) + \frac{79744}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
& = -0.008335007472438759 \dots
\end{aligned} \tag{B.25}$$

$$\begin{aligned}
\langle S_1^x S_3^z S_5^x S_6^z \rangle &= \frac{1}{240} - \frac{11}{20} \zeta_a(1) + \frac{1979}{180} \zeta_a(3) - \frac{833}{15} \zeta_a(1) \zeta_a(3) - \frac{4919}{18} \zeta_a(3)^2 \\
&\quad - \frac{13612}{135} \zeta_a(3)^3 - \frac{75923}{900} \zeta_a(5) + \frac{10660}{9} \zeta_a(1) \zeta_a(5) + \frac{10547}{450} \zeta_a(3) \zeta_a(5) \\
&\quad + \frac{27224}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{41492}{225} \zeta_a(3)^2 \zeta_a(5) - \frac{189064}{45} \zeta_a(5)^2 \\
&\quad + \frac{165968}{45} \zeta_a(1) \zeta_a(5)^2 + \frac{6308}{9} \zeta_a(3) \zeta_a(5)^2 + \frac{9584}{27} \zeta_a(5)^3 + \frac{41321}{180} \zeta_a(7) \\
&\quad - \frac{200018}{45} \zeta_a(1) \zeta_a(7) + \frac{28217}{3} \zeta_a(3) \zeta_a(7) - \frac{290444}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&\quad - \frac{88312}{45} \zeta_a(3)^2 \zeta_a(7) + \frac{59927}{45} \zeta_a(5) \zeta_a(7) - \frac{67088}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&\quad - \frac{162484}{45} \zeta_a(7)^2 + \frac{234808}{45} \zeta_a(1) \zeta_a(7)^2 - \frac{3892}{25} \zeta_a(9) \\
&\quad + 3318 \zeta_a(1) \zeta_a(9) - \frac{665042}{75} \zeta_a(3) \zeta_a(9) + \frac{88312}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&\quad + \frac{67088}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{92848}{15} \zeta_a(5) \zeta_a(9) - \frac{134176}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= 0.0008303788046606665 \dots
\end{aligned} \tag{B.26}$$

$$\begin{aligned}
\langle S_1^x S_3^z S_5^z S_6^x \rangle &= \frac{1}{240} - \frac{43}{60} \zeta_a(1) + \frac{487}{40} \zeta_a(3) - \frac{2599}{45} \zeta_a(1) \zeta_a(3) - \frac{2491}{9} \zeta_a(3)^2 \\
&\quad - \frac{13612}{135} \zeta_a(3)^3 - \frac{156821}{1800} \zeta_a(5) + \frac{3596}{3} \zeta_a(1) \zeta_a(5) + \frac{5161}{225} \zeta_a(3) \zeta_a(5) \\
&\quad + \frac{27224}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{41492}{225} \zeta_a(3)^2 \zeta_a(5) - \frac{189214}{45} \zeta_a(5)^2 \\
&\quad + \frac{165968}{45} \zeta_a(1) \zeta_a(5)^2 + \frac{6308}{9} \zeta_a(3) \zeta_a(5)^2 + \frac{9584}{27} \zeta_a(5)^3 + \frac{10409}{45} \zeta_a(7) \\
&\quad - \frac{200543}{45} \zeta_a(1) \zeta_a(7) + \frac{28238}{3} \zeta_a(3) \zeta_a(7) - \frac{290444}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&\quad - \frac{88312}{45} \zeta_a(3)^2 \zeta_a(7) + \frac{59927}{45} \zeta_a(5) \zeta_a(7) - \frac{67088}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&\quad - \frac{162484}{45} \zeta_a(7)^2 + \frac{234808}{45} \zeta_a(1) \zeta_a(7)^2 - \frac{3892}{25} \zeta_a(9) \\
&\quad + 3318 \zeta_a(1) \zeta_a(9) - \frac{665042}{75} \zeta_a(3) \zeta_a(9) + \frac{88312}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&\quad + \frac{67088}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{92848}{15} \zeta_a(5) \zeta_a(9) - \frac{134176}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= -0.001340720075108033 \dots
\end{aligned} \tag{B.27}$$

$$\begin{aligned}
\langle S_1^x S_2^x S_3^x S_4^x S_5^y S_6^y \rangle &= \frac{1}{2240} + \frac{11}{240} \zeta_a(1) - \frac{23}{16} \zeta_a(3) + \frac{382}{45} \zeta_a(1) \zeta_a(3) + \frac{7961}{180} \zeta_a(3)^2 \\
&+ \frac{148}{9} \zeta_a(3)^3 + \frac{90583}{6300} \zeta_a(5) - \frac{5887}{30} \zeta_a(1) \zeta_a(5) - \frac{4061}{450} \zeta_a(3) \zeta_a(5) \\
&- \frac{296}{3} \zeta_a(1) \zeta_a(3) \zeta_a(5) + \frac{6716}{225} \zeta_a(3)^2 \zeta_a(5) + \frac{430573}{630} \zeta_a(5)^2 \\
&- \frac{26864}{45} \zeta_a(1) \zeta_a(5)^2 - \frac{340}{3} \zeta_a(3) \zeta_a(5)^2 - \frac{402}{7} \zeta_a(5)^3 - \frac{32909}{720} \zeta_a(7) \\
&+ \frac{69223}{90} \zeta_a(1) \zeta_a(7) - \frac{138161}{90} \zeta_a(3) \zeta_a(7) + \frac{47012}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&+ \frac{952}{3} \zeta_a(3)^2 \zeta_a(7) - \frac{39919}{180} \zeta_a(5) \zeta_a(7) + \frac{1206}{5} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&+ \frac{34811}{60} \zeta_a(7)^2 - \frac{4221}{5} \zeta_a(1) \zeta_a(7)^2 + \frac{6543}{200} \zeta_a(9) \\
&- \frac{2908}{5} \zeta_a(1) \zeta_a(9) + \frac{36322}{25} \zeta_a(3) \zeta_a(9) - 952 \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&- \frac{10854}{25} \zeta_a(3)^2 \zeta_a(9) - \frac{4973}{5} \zeta_a(5) \zeta_a(9) + \frac{7236}{5} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= -0.005996024922536831 \dots
\end{aligned} \tag{B.28}$$

$$\begin{aligned}
\langle S_1^x S_2^x S_3^x S_4^y S_5^x S_6^y \rangle &= \frac{1}{2240} - \frac{1}{240} \zeta_a(1) - \frac{643}{1440} \zeta_a(3) + \frac{179}{60} \zeta_a(1) \zeta_a(3) + \frac{1333}{90} \zeta_a(3)^2 \\
&+ \frac{223}{45} \zeta_a(3)^3 + \frac{301507}{50400} \zeta_a(5) - \frac{6139}{90} \zeta_a(1) \zeta_a(5) - \frac{5221}{900} \zeta_a(3) \zeta_a(5) \\
&- \frac{446}{15} \zeta_a(1) \zeta_a(3) \zeta_a(5) + \frac{2018}{225} \zeta_a(3)^2 \zeta_a(5) + \frac{251033}{1260} \zeta_a(5)^2 \\
&- \frac{8072}{45} \zeta_a(1) \zeta_a(5)^2 - 34 \zeta_a(3) \zeta_a(5)^2 - \frac{1084}{63} \zeta_a(5)^3 - \frac{14513}{720} \zeta_a(7) \\
&+ \frac{11753}{45} \zeta_a(1) \zeta_a(7) - \frac{162149}{360} \zeta_a(3) \zeta_a(7) + \frac{14126}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&+ \frac{476}{5} \zeta_a(3)^2 \zeta_a(7) - \frac{24209}{360} \zeta_a(5) \zeta_a(7) + \frac{1084}{15} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&+ \frac{3241}{20} \zeta_a(7)^2 - \frac{3794}{15} \zeta_a(1) \zeta_a(7)^2 + \frac{731}{50} \zeta_a(9) \\
&- \frac{1959}{10} \zeta_a(1) \zeta_a(9) + \frac{21257}{50} \zeta_a(3) \zeta_a(9) - \frac{1428}{5} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&- \frac{3252}{25} \zeta_a(3)^2 \zeta_a(9) - \frac{1389}{5} \zeta_a(5) \zeta_a(9) + \frac{2168}{5} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= 0.001715129839332883 \dots
\end{aligned} \tag{B.29}$$

$$\begin{aligned}
\langle S_1^x S_2^x S_3^x S_4^y S_5^y S_6^x \rangle &= \frac{1}{2240} - \frac{7}{240} \zeta_a(1) - \frac{239}{1440} \zeta_a(3) + \frac{86}{45} \zeta_a(1) \zeta_a(3) + \frac{1123}{90} \zeta_a(3)^2 \\
&+ \frac{223}{45} \zeta_a(3)^3 + \frac{82489}{16800} \zeta_a(5) - \frac{5309}{90} \zeta_a(1) \zeta_a(5) - \frac{2843}{450} \zeta_a(3) \zeta_a(5) \\
&- \frac{446}{15} \zeta_a(1) \zeta_a(3) \zeta_a(5) + \frac{2018}{225} \zeta_a(3)^2 \zeta_a(5) + \frac{247883}{1260} \zeta_a(5)^2 \\
&- \frac{8072}{45} \zeta_a(1) \zeta_a(5)^2 - 34 \zeta_a(3) \zeta_a(5)^2 - \frac{1084}{63} \zeta_a(5)^3 - \frac{2785}{144} \zeta_a(7) \\
&+ \frac{45563}{180} \zeta_a(1) \zeta_a(7) - \frac{160259}{360} \zeta_a(3) \zeta_a(7) + \frac{14126}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&+ \frac{476}{5} \zeta_a(3)^2 \zeta_a(7) - \frac{24209}{360} \zeta_a(5) \zeta_a(7) + \frac{1084}{15} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&+ \frac{3241}{20} \zeta_a(7)^2 - \frac{3794}{15} \zeta_a(1) \zeta_a(7)^2 + \frac{731}{50} \zeta_a(9) \\
&- \frac{1959}{10} \zeta_a(1) \zeta_a(9) + \frac{21257}{50} \zeta_a(3) \zeta_a(9) - \frac{1428}{5} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&- \frac{3252}{25} \zeta_a(3)^2 \zeta_a(9) - \frac{1389}{5} \zeta_a(5) \zeta_a(9) + \frac{2168}{5} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= -0.002589603677721155 \dots
\end{aligned} \tag{B.30}$$

$$\begin{aligned}
\langle S_1^x S_2^x S_3^y S_4^x S_5^x S_6^y \rangle &= \frac{1}{2240} - \frac{11}{240} \zeta_a(1) + \frac{1537}{1440} \zeta_a(3) - \frac{421}{60} \zeta_a(1) \zeta_a(3) - \frac{317}{8} \zeta_a(3)^2 \\
&- \frac{79}{5} \zeta_a(3)^3 - \frac{623431}{50400} \zeta_a(5) + \frac{7949}{45} \zeta_a(1) \zeta_a(5) + \frac{15851}{1800} \zeta_a(3) \zeta_a(5) \\
&+ \frac{474}{5} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{6404}{225} \zeta_a(3)^2 \zeta_a(5) - \frac{808669}{1260} \zeta_a(5)^2 \\
&+ \frac{25616}{45} \zeta_a(1) \zeta_a(5)^2 + 108 \zeta_a(3) \zeta_a(5)^2 + \frac{3452}{63} \zeta_a(5)^3 + \frac{15553}{360} \zeta_a(7) \\
&- \frac{64771}{90} \zeta_a(1) \zeta_a(7) + \frac{172993}{120} \zeta_a(3) \zeta_a(7) - \frac{44828}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&- \frac{1512}{5} \zeta_a(3)^2 \zeta_a(7) + \frac{75247}{360} \zeta_a(5) \zeta_a(7) - \frac{3452}{15} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&- \frac{32809}{60} \zeta_a(7)^2 + \frac{12082}{15} \zeta_a(1) \zeta_a(7)^2 - \frac{1593}{50} \zeta_a(9) \\
&+ \frac{5503}{10} \zeta_a(1) \zeta_a(9) - \frac{68441}{50} \zeta_a(3) \zeta_a(9) + \frac{4536}{5} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&+ \frac{10356}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{4687}{5} \zeta_a(5) \zeta_a(9) - \frac{6904}{5} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= -0.002089292530622660 \dots
\end{aligned} \tag{B.31}$$

$$\begin{aligned}
\langle S_1^x S_2^y S_3^x S_4^x S_5^x S_6^y \rangle &= \frac{1}{2240} - \frac{19}{240} \zeta_a(1) + \frac{95}{48} \zeta_a(3) - \frac{2087}{180} \zeta_a(1) \zeta_a(3) - \frac{22057}{360} \zeta_a(3)^2 \\
&\quad - \frac{3134}{135} \zeta_a(3)^3 - \frac{12221}{630} \zeta_a(5) + \frac{2707}{10} \zeta_a(1) \zeta_a(5) + \frac{3971}{360} \zeta_a(3) \zeta_a(5) \\
&\quad + \frac{6268}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{1897}{45} \zeta_a(3)^2 \zeta_a(5) - \frac{121259}{126} \zeta_a(5)^2 \\
&\quad + \frac{7588}{9} \zeta_a(1) \zeta_a(5)^2 + \frac{1441}{9} \zeta_a(3) \zeta_a(5)^2 + \frac{15340}{189} \zeta_a(5)^3 + \frac{44329}{720} \zeta_a(7) \\
&\quad - \frac{191611}{180} \zeta_a(1) \zeta_a(7) + \frac{194381}{90} \zeta_a(3) \zeta_a(7) - \frac{13279}{9} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&\quad - \frac{20174}{45} \zeta_a(3)^2 \zeta_a(7) + \frac{2798}{9} \zeta_a(5) \zeta_a(7) - \frac{3068}{9} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&\quad - \frac{14777}{18} \zeta_a(7)^2 + \frac{10738}{9} \zeta_a(1) \zeta_a(7)^2 - \frac{1763}{40} \zeta_a(9) \\
&\quad + \frac{4029}{5} \zeta_a(1) \zeta_a(9) - \frac{12269}{6} \zeta_a(3) \zeta_a(9) + \frac{20174}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&\quad + \frac{3068}{5} \zeta_a(3)^2 \zeta_a(9) + \frac{4222}{3} \zeta_a(5) \zeta_a(9) - \frac{6136}{3} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= 0.0008959339234083256 \dots
\end{aligned} \tag{B.32}$$

$$\begin{aligned}
\langle S_1^x S_2^x S_3^y S_4^x S_5^y S_6^x \rangle &= \frac{1}{2240} - \frac{13}{240} \zeta_a(1) + \frac{619}{480} \zeta_a(3) - \frac{229}{30} \zeta_a(1) \zeta_a(3) - \frac{14687}{360} \zeta_a(3)^2 \\
&\quad - \frac{79}{5} \zeta_a(3)^3 - \frac{74279}{5600} \zeta_a(5) + \frac{2723}{15} \zeta_a(1) \zeta_a(5) + \frac{5197}{600} \zeta_a(3) \zeta_a(5) \\
&\quad + \frac{474}{5} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{6404}{225} \zeta_a(3)^2 \zeta_a(5) - \frac{810139}{1260} \zeta_a(5)^2 \\
&\quad + \frac{25616}{45} \zeta_a(1) \zeta_a(5)^2 + 108 \zeta_a(3) \zeta_a(5)^2 + \frac{3452}{63} \zeta_a(5)^3 + \frac{2633}{60} \zeta_a(7) \\
&\quad - \frac{8687}{12} \zeta_a(1) \zeta_a(7) + \frac{173287}{120} \zeta_a(3) \zeta_a(7) - \frac{44828}{45} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&\quad - \frac{1512}{5} \zeta_a(3)^2 \zeta_a(7) + \frac{75247}{360} \zeta_a(5) \zeta_a(7) - \frac{3452}{15} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&\quad - \frac{32809}{60} \zeta_a(7)^2 + \frac{12082}{15} \zeta_a(1) \zeta_a(7)^2 - \frac{1593}{50} \zeta_a(9) \\
&\quad + \frac{5503}{10} \zeta_a(1) \zeta_a(9) - \frac{68441}{50} \zeta_a(3) \zeta_a(9) + \frac{4536}{5} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&\quad + \frac{10356}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{4687}{5} \zeta_a(5) \zeta_a(9) - \frac{6904}{5} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= 0.001508610170462532 \dots
\end{aligned} \tag{B.33}$$

$$\begin{aligned}
\langle S_1^x S_2^y S_3^x S_4^z S_5^z S_6^y \rangle &= \frac{1}{6720} - \frac{1}{80}\zeta_a(1) + \frac{25}{288}\zeta_a(3) - \frac{16}{45}\zeta_a(1)\zeta_a(3) - \frac{121}{90}\zeta_a(3)^2 \\
&\quad - \frac{73}{135}\zeta_a(3)^3 + \frac{1541}{50400}\zeta_a(5) + \frac{413}{90}\zeta_a(1)\zeta_a(5) - \frac{539}{450}\zeta_a(3)\zeta_a(5) \\
&\quad + \frac{146}{45}\zeta_a(1)\zeta_a(3)\zeta_a(5) - \frac{211}{225}\zeta_a(3)^2\zeta_a(5) - \frac{32891}{1260}\zeta_a(5)^2 \\
&\quad + \frac{844}{45}\zeta_a(1)\zeta_a(5)^2 + \frac{32}{9}\zeta_a(3)\zeta_a(5)^2 + \frac{344}{189}\zeta_a(5)^3 - \frac{137}{120}\zeta_a(7) \\
&\quad - \frac{217}{15}\zeta_a(1)\zeta_a(7) + \frac{2578}{45}\zeta_a(3)\zeta_a(7) - \frac{1477}{45}\zeta_a(1)\zeta_a(3)\zeta_a(7) \\
&\quad - \frac{448}{45}\zeta_a(3)^2\zeta_a(7) + \frac{107}{15}\zeta_a(5)\zeta_a(7) - \frac{344}{45}\zeta_a(3)\zeta_a(5)\zeta_a(7) \\
&\quad - \frac{2219}{90}\zeta_a(7)^2 + \frac{1204}{45}\zeta_a(1)\zeta_a(7)^2 + \frac{207}{200}\zeta_a(9) \\
&\quad + \frac{103}{10}\zeta_a(1)\zeta_a(9) - \frac{8017}{150}\zeta_a(3)\zeta_a(9) + \frac{448}{15}\zeta_a(1)\zeta_a(3)\zeta_a(9) \\
&\quad + \frac{344}{25}\zeta_a(3)^2\zeta_a(9) + \frac{634}{15}\zeta_a(5)\zeta_a(9) - \frac{688}{15}\zeta_a(1)\zeta_a(5)\zeta_a(9) \\
&= 0.0007223711440396728 \dots
\end{aligned} \tag{B.34}$$

$$\begin{aligned}
\langle S_1^x S_2^x S_3^y S_4^z S_5^y S_6^z \rangle &= \frac{1}{6720} + \frac{1}{80}\zeta_a(1) - \frac{353}{720}\zeta_a(3) + \frac{569}{180}\zeta_a(1)\zeta_a(3) + \frac{327}{20}\zeta_a(3)^2 \\
&\quad + \frac{163}{27}\zeta_a(3)^3 + \frac{26927}{5040}\zeta_a(5) - \frac{655}{9}\zeta_a(1)\zeta_a(5) - \frac{109}{30}\zeta_a(3)\zeta_a(5) \\
&\quad - \frac{326}{9}\zeta_a(1)\zeta_a(3)\zeta_a(5) + \frac{488}{45}\zeta_a(3)^2\zeta_a(5) + \frac{63055}{252}\zeta_a(5)^2 \\
&\quad - \frac{1952}{9}\zeta_a(1)\zeta_a(5)^2 - \frac{370}{9}\zeta_a(3)\zeta_a(5)^2 - \frac{3940}{189}\zeta_a(5)^3 - \frac{209}{12}\zeta_a(7) \\
&\quad + \frac{4291}{15}\zeta_a(1)\zeta_a(7) - \frac{22497}{40}\zeta_a(3)\zeta_a(7) + \frac{3416}{9}\zeta_a(1)\zeta_a(3)\zeta_a(7) \\
&\quad + \frac{1036}{9}\zeta_a(3)^2\zeta_a(7) - \frac{5869}{72}\zeta_a(5)\zeta_a(7) + \frac{788}{9}\zeta_a(3)\zeta_a(5)\zeta_a(7) \\
&\quad + \frac{7609}{36}\zeta_a(7)^2 - \frac{2758}{9}\zeta_a(1)\zeta_a(7)^2 + \frac{251}{20}\zeta_a(9) \\
&\quad - \frac{433}{2}\zeta_a(1)\zeta_a(9) + \frac{15961}{30}\zeta_a(3)\zeta_a(9) - \frac{1036}{3}\zeta_a(1)\zeta_a(3)\zeta_a(9) \\
&\quad - \frac{788}{5}\zeta_a(3)^2\zeta_a(9) - \frac{1087}{3}\zeta_a(5)\zeta_a(9) + \frac{1576}{3}\zeta_a(1)\zeta_a(5)\zeta_a(9) \\
&= 0.001419281470187951 \dots
\end{aligned} \tag{B.35}$$

$$\begin{aligned}
\langle S_1^x S_2^x S_3^y S_4^z S_5^z S_6^y \rangle &= \frac{1}{6720} + \frac{1}{120} \zeta_a(1) - \frac{11}{24} \zeta_a(3) + \frac{571}{180} \zeta_a(1) \zeta_a(3) + \frac{733}{45} \zeta_a(3)^2 \\
&+ \frac{163}{27} \zeta_a(3)^3 + \frac{26003}{5040} \zeta_a(5) - \frac{145}{2} \zeta_a(1) \zeta_a(5) - \frac{73}{20} \zeta_a(3) \zeta_a(5) \\
&- \frac{326}{9} \zeta_a(1) \zeta_a(3) \zeta_a(5) + \frac{488}{45} \zeta_a(3)^2 \zeta_a(5) + \frac{63041}{252} \zeta_a(5)^2 \\
&- \frac{1952}{9} \zeta_a(1) \zeta_a(5)^2 - \frac{370}{9} \zeta_a(3) \zeta_a(5)^2 - \frac{3940}{189} \zeta_a(5)^3 - \frac{3107}{180} \zeta_a(7) \\
&+ \frac{51443}{180} \zeta_a(1) \zeta_a(7) - \frac{67477}{120} \zeta_a(3) \zeta_a(7) + \frac{3416}{9} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&+ \frac{1036}{9} \zeta_a(3)^2 \zeta_a(7) - \frac{5869}{72} \zeta_a(5) \zeta_a(7) + \frac{788}{9} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&+ \frac{7609}{36} \zeta_a(7)^2 - \frac{2758}{9} \zeta_a(1) \zeta_a(7)^2 + \frac{251}{20} \zeta_a(9) \\
&- \frac{433}{2} \zeta_a(1) \zeta_a(9) + \frac{15961}{30} \zeta_a(3) \zeta_a(9) - \frac{1036}{3} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&- \frac{788}{5} \zeta_a(3)^2 \zeta_a(9) - \frac{1087}{3} \zeta_a(5) \zeta_a(9) + \frac{1576}{3} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= -0.002041037236271903 \dots
\end{aligned} \tag{B.36}$$

$$\begin{aligned}
\langle S_1^x S_2^y S_3^z S_4^x S_5^z S_6^y \rangle &= \frac{1}{6720} - \frac{7}{240} \zeta_a(1) + \frac{1169}{1440} \zeta_a(3) - \frac{941}{180} \zeta_a(1) \zeta_a(3) - \frac{3391}{120} \zeta_a(3)^2 \\
&- \frac{1474}{135} \zeta_a(3)^3 - \frac{150971}{16800} \zeta_a(5) + \frac{5662}{45} \zeta_a(1) \zeta_a(5) + \frac{1237}{200} \zeta_a(3) \zeta_a(5) \\
&+ \frac{2948}{45} \zeta_a(1) \zeta_a(3) \zeta_a(5) - \frac{1474}{75} \zeta_a(3)^2 \zeta_a(5) - \frac{281161}{630} \zeta_a(5)^2 \\
&+ \frac{5896}{15} \zeta_a(1) \zeta_a(5)^2 + \frac{671}{9} \zeta_a(3) \zeta_a(5)^2 + \frac{7148}{189} \zeta_a(5)^3 + \frac{21893}{720} \zeta_a(7) \\
&- \frac{22673}{45} \zeta_a(1) \zeta_a(7) + \frac{5013}{5} \zeta_a(3) \zeta_a(7) - \frac{10318}{15} \zeta_a(1) \zeta_a(3) \zeta_a(7) \\
&- \frac{9394}{45} \zeta_a(3)^2 \zeta_a(7) + \frac{2179}{15} \zeta_a(5) \zeta_a(7) - \frac{7148}{45} \zeta_a(3) \zeta_a(5) \zeta_a(7) \\
&- \frac{17059}{45} \zeta_a(7)^2 + \frac{25018}{45} \zeta_a(1) \zeta_a(7)^2 - \frac{4441}{200} \zeta_a(9) \\
&+ \frac{1917}{5} \zeta_a(1) \zeta_a(9) - \frac{71282}{75} \zeta_a(3) \zeta_a(9) + \frac{9394}{15} \zeta_a(1) \zeta_a(3) \zeta_a(9) \\
&+ \frac{7148}{25} \zeta_a(3)^2 \zeta_a(9) + \frac{9748}{15} \zeta_a(5) \zeta_a(9) - \frac{14296}{15} \zeta_a(1) \zeta_a(5) \zeta_a(9) \\
&= -0.0001528594146694370 \dots
\end{aligned} \tag{B.37}$$

$$\begin{aligned}
\langle S_1^x S_2^y S_3^z S_4^z S_5^y S_6^x \rangle &= \frac{1}{6720} - \frac{1}{24}\zeta_a(1) + \frac{41}{36}\zeta_a(3) - \frac{109}{18}\zeta_a(1)\zeta_a(3) - \frac{571}{18}\zeta_a(3)^2 \\
&\quad - \frac{529}{45}\zeta_a(3)^3 - \frac{134587}{12600}\zeta_a(5) + \frac{6328}{45}\zeta_a(1)\zeta_a(5) + \frac{2737}{450}\zeta_a(3)\zeta_a(5) \\
&\quad + \frac{1058}{15}\zeta_a(1)\zeta_a(3)\zeta_a(5) - \frac{4852}{225}\zeta_a(3)^2\zeta_a(5) - \frac{617377}{1260}\zeta_a(5)^2 \\
&\quad + \frac{19408}{45}\zeta_a(1)\zeta_a(5)^2 + 82\zeta_a(3)\zeta_a(5)^2 + \frac{872}{21}\zeta_a(5)^3 + \frac{2599}{80}\zeta_a(7) \\
&\quad - \frac{8197}{15}\zeta_a(1)\zeta_a(7) + \frac{10999}{10}\zeta_a(3)\zeta_a(7) - \frac{33964}{45}\zeta_a(1)\zeta_a(3)\zeta_a(7) \\
&\quad - \frac{1148}{5}\zeta_a(3)^2\zeta_a(7) + \frac{7132}{45}\zeta_a(5)\zeta_a(7) - \frac{872}{5}\zeta_a(3)\zeta_a(5)\zeta_a(7) \\
&\quad - \frac{2086}{5}\zeta_a(7)^2 + \frac{3052}{5}\zeta_a(1)\zeta_a(7)^2 - \frac{4581}{200}\zeta_a(9) \\
&\quad + \frac{4121}{10}\zeta_a(1)\zeta_a(9) - \frac{26024}{25}\zeta_a(3)\zeta_a(9) + \frac{3444}{5}\zeta_a(1)\zeta_a(3)\zeta_a(9) \\
&\quad + \frac{7848}{25}\zeta_a(3)^2\zeta_a(9) + \frac{3576}{5}\zeta_a(5)\zeta_a(9) - \frac{5232}{5}\zeta_a(1)\zeta_a(5)\zeta_a(9) \\
&= -0.0003767104281403536 \dots
\end{aligned} \tag{B.38}$$

References

- [1] H.A. Bethe, Z. Phys. **71** (1931) 205.
- [2] M. Takahashi, *Thermodynamics of One-Dimensional Solvable Models*, Cambridge University Press, Cambridge, 1999.
- [3] E. Lieb, T. Schultz, D. Mattis, Ann. Phys. (N.Y.) **16** (1961) 407.
- [4] B.M. McCoy, Phys. Rev. **173** (1968) 531.
- [5] L. Hulthén, Arkiv Mat. Astron. Fysik **A 26** (1938) 1.
- [6] M. Takahashi, J. Phys. C: Solid State Phys. **10** (1977) 1289.
- [7] J. Dittrich, V.I. Inozemtsev, J. Phys. A: Math. Gen. **30** (1997) L623.
- [8] M. Jimbo, K. Miki, T. Miwa, A. Nakayashiki, Phys. Lett. A **168** (1992) 256.
- [9] M. Jimbo, T. Miwa, *Algebraic Analysis of Solvable Lattice Models*, CBMS Regional Conference Series in Mathematics vol.**85**, American Mathematical Society, Providence, 1994.
- [10] A. Nakayashiki, Int. J. Mod. Phys. A **9** (1994) 5673.
- [11] V.E. Korepin, A. Izergin, F.H.L. Essler, D. Uglov, Phys. Lett. A **190** (1994) 182.
- [12] M. Jimbo, T. Miwa, J. Phys. A: Math. Gen. **29** (1996) 2923.

- [13] N. Kitanine, J.M. Maillet, V. Terras, Nucl. Phys. B **567** (2000), 554.
- [14] F.Göhmman, A. Klümper, A. Seel, J. Phys. A: Math. Gen **37** (2004) 7625.
- [15] F.Göhmman, A. Klümper, A. Seel, J. Phys. A: Math. Gen **38** (2005) 1833.
- [16] H.E. Boos, V.E. Korepin, J. Phys. A: Math. Gen. **34** (2001) 5311.
- [17] H.E. Boos, V.E. Korepin, “*Evaluation of integrals representing correlators in XXX Heisenberg spin chain*” in. MathPhys Odyssey 2001, Birkhäuser, Basel, (2001) 65.
- [18] H.E. Boos, V.E. Korepin, Y. Nishiyama, M. Shiroishi, J. Phys. A: Math. Gen **35** (2002) 4443.
- [19] K. Sakai, M. Shiroishi, Y. Nishiyama, M. Takahashi, Phys. Rev. E **67** (2003) 065101.
- [20] G. Kato, M. Shiroishi, M. Takahashi, K. Sakai, J. Phys. A: Math. Gen. **36** (2003) L337.
- [21] M. Takahashi, G. Kato, M. Shiroishi, J. Phys. Soc. Jpn, **73** (2004) 245.
- [22] G. Kato, M. Shiroishi, M. Takahashi, K. Sakai, J. Phys. A: Math. Gen. **37** (2004) 5097.
- [23] H.E. Boos, V.E. Korepin, F.A. Smirnov, Nucl. Phys. B **658** (2003) 417.
- [24] V.G. Knizhnik, A.B. Zamolodchikov, Nucl. Phys. B **247** (1984) 83.
- [25] I. Frenkel, N. Reshetikhin, Comm. Math. Phys. **146** (1992) 1.
- [26] F.A. Smirnov, *Form Factors in Completely Integrable Models of Quantum Field Theory*, Adv. Series in Math. Phys. 14, World Scientific, Singapore (1992).
- [27] F.A. Smirnov, Int. Jour. Math. Phys. (Suppl. 1B), A7 (1992) 813.
- [28] H.E. Boos, M. Jimbo, T. Miwa, F. Smirnov, Y. Takeyama, Algebra Anal. **17** (2005) 115.
- [29] H.E. Boos, M. Jimbo, T. Miwa, F. Smirnov, Y. Takeyama, Commun. Math. Phys. **261** (2006) 245.
- [30] H.E. Boos, M. Jimbo, T. Miwa, F. Smirnov, Y. Takeyama, J. Phys. A: Math. Gen. **38** (2005) 7629.
- [31] H.E. Boos, M. Jimbo, T. Miwa, F. Smirnov, Y. Takeyama, Lett. Math. Phys. **75** (2006) 201.
- [32] H.E. Boos, M. Jimbo, T. Miwa, F. Smirnov, Y. Takeyama, “*Algebraic representation of correlation functions in integrable spin chains*”, hep-th/0601132.
- [33] H.E. Boos, M. Jimbo, T. Miwa, F. Smirnov, Y. Takeyama, “*Hidden Grassmann structure in the XXZ model*”, hep-th/0606280.
- [34] H.E. Boos, M. Shiroishi, M. Takahashi, Nucl. Phys. B **712** (2005) 573.
- [35] J. Sato, M. Shiroishi, J. Phys. A: Math. Gen. **38** (2005) L405.

- [36] J. Sato, M. Shiroishi, M. Takahashi, Nucl. Phys. B **729** (2005) 441, hep-th/0507290.
- [37] M. Bortz, J. Sato, M. Shiroishi, in preparation.
- [38] N. Muramoto, M. Takahashi, J. Phys. Soc. Jpn. **68** (1999) 2098.
- [39] N. Kawakami, S.K. Yang, “*Conformal Field Theory and One-Dimensional Quantum Systems*”, (Iwanami, Tokyo, 1997) (in Japanese).
- [40] V.E. Korepin, S. Lukyanov, Y. Nishiyama, M. Shiroishi, Phys. Lett. A **312** (2003) 21.
- [41] N. Kitanine, J.M. Maillet, N.A. Slavnov, V. Terras, J. Phys. A: Math. Gen. **35** (2002) L753.
- [42] Y. Nishiyama, in private communication.
- [43] G. Vidal, J.I. Latorre, E. Rico, A. Kitaev, Phys. Rev. Lett. **90** (2003) 227902.
- [44] C. Holzhey, F. Larsen, F. Wilczek, Nucl. Phys. B **424** (1994) 443.